产品名称: CGGBP1 Rabbit Polyclonal Antibody

产品货号: APRab08720



产品概述 (Summary)

产品名称 (Production Name) CGGBP1 Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application)IHC,ICC/IF,ELISA种属反应性 (Reactivity)Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) CGGBP1

CGGBP1; CGGBP; CGG triplet repeat-binding protein 1; CGG-binding protein 1; **别名 (Alternative Names)**

20 kDa CGG-binding protein; p20-CGGBP DNA-binding protein

基因 ID (Gene ID) 8545.0

Q9UFW8.The antiserum was produced against synthesized peptide derived 蛋白ID (SwissProt ID)

from human CGGBP1. AA range:41-90

产品应用 (Application)

稀释比 (Dilution Ratio) IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:10000

蛋白分子量 (Molecular Weight)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: CGGBP1 Rabbit Polyclonal Antibody

产品货号: APRab08720




研究背景 (Background)

CGGBP1 influences expression of the FMR1 gene (MIM 309550), which is associated with the fragile X mental retardation syndrome (MIM 300624), by specifically interacting with the 5-prime (CGG)n-3-prime repeat in its 5-prime UTR.[supplied by OMIM, Mar 2008], developmental stage: Expressed in fetal brain and kidney. Lower expression in fetal liver and lung., function: Binds to nonmethylated 5'-d(CGG)(n)-3' trinucleotide repeats in the FMR1 promoter. May play a role in regulating FMR1 promoter., miscellaneous: Binding is severely inhibited by complete or partial cytosine-specific DNA methylation of the binding motif., PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., tissue specificity: Ubiquitous. Highly expressed in placenta, thymus, lymph nodes, cerebellum and cerebral cortex. Low expression in other regions of the brain.,

研究领域 (Research Area)

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200 (4° overnight) . 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min) .

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838