

产品名称: CKR-5 Rabbit Polyclonal Antibody  
产品货号: APRab08871



## 产品概述 (Summary)

|                        |                                  |
|------------------------|----------------------------------|
| 产品名称 (Production Name) | CKR-5 Rabbit Polyclonal Antibody |
| 描述 (Description)       | Rabbit polyclonal Antibody       |
| 宿主 (Host)              | Rabbit                           |
| 应用 (Application)       | IHC, ICC, IF, ELISA              |
| 种属反应性 (Reactivity)     | Human, Mouse, Rat                |

## 产品性能 (Performance)

|                     |                                                                                                   |
|---------------------|---------------------------------------------------------------------------------------------------|
| 偶联物 (Conjugation)   | Unconjugated                                                                                      |
| 修饰 (Modification)   | Unmodified                                                                                        |
| 同种型 (Isotype)       | IgG                                                                                               |
| 克隆 (Clonality)      | Polyclonal                                                                                        |
| 形式 (Form)           | Liquid                                                                                            |
| 存放说明 (Storage)      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.          |
| 储存溶液 (Buffer)       | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| 纯化方式 (Purification) | Affinity purification                                                                             |

## 免疫原信息 (Immunogen)

|                        |                                                                                                                                   |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------|
| 基因名 (Gene Name)        | CCR5                                                                                                                              |
| 别名 (Alternative Names) | CCR5; CMKBR5; C-C chemokine receptor type 5; C-C CKR-5; CC-CKR-5; CCR-5; CCR5; CHEMR13; HIV-1 fusion coreceptor; CD antigen CD195 |
| 基因 ID (Gene ID)        | 1234.0                                                                                                                            |
| 蛋白 ID (SwissProt ID)   | P51681. The antiserum was produced against synthesized peptide derived from human CCR5. AA range: 292-341                         |

## 产品应用 (Application)

|                          |                                                          |
|--------------------------|----------------------------------------------------------|
| 稀释比 (Dilution Ratio)     | IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:5000-1:10000 |
| 蛋白分子量 (Molecular Weight) | 41kDa                                                    |

**产品名称: CKR-5 Rabbit Polyclonal Antibody**  
**产品货号: APRab08871**



## 研究背景 (Background)

This gene encodes a member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. This protein is expressed by T cells and macrophages, and is known to be an important co-receptor for macrophage-tropic virus, including HIV, to enter host cells. Defective alleles of this gene have been associated with the HIV infection resistance. The ligands of this receptor include monocyte chemoattractant protein 2 (MCP-2), macrophage inflammatory protein 1 alpha (MIP-1 alpha), macrophage inflammatory protein 1 beta (MIP-1 beta) and regulated on activation normal T expressed and secreted protein (RANTES). Expression of this gene was also detected in a promyeloblastic cell line, suggesting that this protein may play a role in granulocyte lineage proliferation and differentiation. This gene is located at the chemokine disease: Genetic variation in CCR5 is associated with susceptibility to insulin-dependent diabetes mellitus type 2 (IDDM2) [MIM:612522]. IDDM is caused by the body's own immune system which destroys the insulin-producing beta cells in the pancreas. Classical features are polydipsia, polyphagia and polyuria, due to hyperglycemia-induced osmotic diuresis., function: Receptor for a number of inflammatory CC-chemokines including MIP-1-alpha, MIP-1-beta and RANTES and subsequently transduces a signal by increasing the intracellular calcium ion level. May play a role in the control of granulocytic lineage proliferation or differentiation. Acts as a coreceptor (CD4 being the primary receptor) for HIV-1 R5 isolates., online information: CC chemokine receptors entry, online information: CCR5 receptor entry, polymorphism: Ser-60 variant, a naturally occurring mutation in a conserved residue in the first intracellular domain of CCR5, results in reduced amounts of the protein in the membrane and consequently may be associated with reduced susceptibility to infection by microbes that depend on these molecules as their receptors., polymorphism: Variations in CCR5 are associated with resistance or susceptibility to immunodeficiency virus type 1 (resistance or susceptibility to HIV-1) [MIM:609423]. Variations in CCR5 gene also influence the rate of progression to AIDS after infection., polymorphism: Variations in CCR5 are associated with susceptibility to West Nile virus (WNV) infection [MIM:610379]., PTM: O-glycosylated, but not N-glycosylated. Ser-6 appears to be the major site. Also sialylated glycans present which contribute to chemokine binding. Thr-16 and Ser-17 may also be glycosylated and, if so, with small moieties such as a T-antigen., PTM: Palmitoylation in the C-terminal is important for cell surface expression, and to a lesser extent, for HIV entry., PTM: Phosphorylation on serine residues in the C-terminal is stimulated by binding CC chemokines especially by APO-RANTES., PTM: Sulfated on at least 2 of the N-terminal tyrosines. Sulfation contributes to the efficiency of HIV-1 entry and is required for efficient binding of the chemokines, CCL3 and CCL4., similarity: Belongs to the G-protein coupled receptor 1 family., subunit: Interacts with PRAF2. Interacts with HIV-1 surface protein gp120. Efficient ligand binding to CCL3/MIP-1alpha and CCR4/MIP-1beta requires sulfation, O-glycosylation and sialic acid modifications. Glycosylation on Ser-6 is required for efficient binding of CCL4. Interacts with ADRBK1., tissue specificity: Highly expressed in spleen, thymus, in the myeloid cell line THP-1, in the promyeloblastic cell line KG-1A and on CD4+ and CD8+ T-cells. Medium levels in peripheral blood leukocytes and in small intestine. Low levels in ovary and lung.,


## 研究领域 (Research Area)

Cytokine-cytokine receptor interaction; Chemokine; Endocytosis;

产品名称: CKR-5 Rabbit Polyclonal Antibody  
产品货号: APRab08871



## 图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using CCR5 Antibody. The picture on the right is blocked with the synthesized peptide.

## 注意事项 (Note)

For research use only .