

产品名称: Cleaved-GGT4 HC (T472) Rabbit Polyclonal Antibody
产品货号: APRab08993



产品概述 (Summary)

产品名称 (Production Name)	Cleaved-GGT4 HC (T472) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	GGT7
别名 (Alternative Names)	GGT7; GGTL3; GGTL5; Gamma-glutamyltransferase 7; GGT 7; Gamma-glutamyltransferase-like 3; Gamma-glutamyltransferase-like 5; Gamma-glutamyltranspeptidase 7; Glutathione hydrolase 7
基因 ID (Gene ID)	2686.0
蛋白 ID (SwissProt ID)	Q9UJ14.The antiserum was produced against synthesized peptide derived from human Gamma-glutamyltransferase 4. AA range:423-472

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:5000-1:20000
----------------------	--------------------------------------

产品名称: Cleaved-GGT4 HC (T472) Rabbit Polyclonal Antibody
产品货号: APRab08993



蛋白分子量 (Molecular Weight) 50kDa


研究背景 (Background)

This gene is a member of a gene family that encodes enzymes involved in both the metabolism of glutathione and in the transpeptidation of amino acids. Changes in the activity of gamma-glutamyltransferase may signal preneoplastic or toxic conditions in the liver or kidney. The protein encoded by this gene consists of a heavy and a light chain, and it can interact with CT120, a plasma membrane-associated protein that is possibly involved in lung carcinogenesis. [provided by RefSeq, Jul 2008],catalytic activity:(5-L-glutamyl)-peptide + an amino acid = peptide + 5-L-glutamyl amino acid.,function:Cleaves glutathione conjugates.,pathway:Sulfur metabolism; glutathione metabolism.,similarity:Belongs to the gamma-glutamyltransferase family.,subunit:Heterodimer composed of the light and heavy chains. The active site is located in the light chain (By similarity). Isoform 3 interacts with FAM57A.,tissue specificity:Widely expressed, but at low level, except in the airway epithelial cells. Detected in brain, heart, kidney, liver, lung, spleen, testis and trachea.,

研究领域 (Research Area)

Taurine and hypotaurine metabolism;Selenoamino acid metabolism;Cyanoamino acid metabolism;Glutathione metabolism;Arachidonic acid metabolism;

图片 (Image Data)



Western blot analysis of lysates from Jurkat cells, treated with etoposide 25uM 24h, using Gamma-glutamyltransferase 4 (heavy chain,Cleaved-Thr472) Antibody. The lane on the right is blocked with the synthesized peptide.

注意事项 (Note)

For research use only .