



产品货号: APRab09008



#### 产品概述 (Summary)

Cleaved-MMP-1 22k (F100) Rabbit Polyclonal Antibody 产品名称 (Production Name)

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit 应用 (Application) WB.ELISA

种属反应性 (Reactivity) Human, Rat, Mouse

#### 产品性能 (Performance)

Unconjugated 偶联物 (Conjugation) 修饰 (Modification) Unmodified

同种型 (Isotype) IgG

Polyclonal 克隆 (Clonality) 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

# 免疫原信息 (Immunogen)

基因名 (Gene Name) MMP1

MMP1; CLG; Interstitial collagenase; Fibroblast collagenase; Matrix 别名 (Alternative Names)

metalloproteinase-1; MMP-1

基因 ID (Gene ID) 4312.0

P03956.The antiserum was produced against synthesized peptide derived 蛋白 ID (SwissProt ID)

from human MMP1. AA range:81-130

## 产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,ELISA 1:5000-1:20000

蛋白分子量 (Molecular Weight) 19kDa

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技支持) Web: https://www.enkilife.cn Tel: 027-87002838

**Antibody** 

产品货号: APRab09008



#### 研究背景 (Background)

matrix metallopeptidase 1(MMP1) Homo sapiens This gene encodes a member of the peptidase M10 family of matrix metalloproteinases (MMPs). Proteins in this family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. The encoded preproprotein is proteolytically processed to generate the mature protease. This secreted protease breaks down the interstitial collagens, including types I, II, and III. The gene is part of a cluster of MMP genes on chromosome 11. Mutations in this gene are associated with chronic obstructive pulmonary disease (COPD). Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Jan 2016], catalytic activity: Cleavage of the triple helix of collagen at about three-guarters of the length of the molecule from the N-terminus, at 775-Gly-|-lle-776 in the alpha-1(I) chain. Cleaves synthetic substrates and alpha-macroglobulins at bonds where P1' is a hydrophobic residue., cofactor: Binds 2 zinc ions per subunit., cofactor: Binds 4 calcium ions per subunit., domain: The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme, domain: There are two distinct domains in this protein; the catalytic Nterminal, and the C-terminal which is involved in substrate specificity and in binding TIMP (tissue inhibitor of metalloproteinases), enzyme regulation: Can be activated without removal of the activation peptide, function: Cleaves collagens of types I, II, and III at one site in the helical domain. Also cleaves collagens of types VII and X. In case of HIV infection, interacts and cleaves the secreted viral Tat protein, leading to a decrease in neuronal Tat's mediated neurotoxicity.,online information:Collagenase entry,PTM:Undergoes autolytic cleavage to two major forms (22 kDa and 27 kDa). A minor form (25 kDa) is the glycosylated form of the 22 kDa form. The 27 kDa form has no activity while the 22/25 kDa form can act as activator for collagenase., similarity: Belongs to the peptidase M10A family., similarity: Contains 4 hemopexin-like domains., subunit: Interacts with HIV-1 Tat.,

## 研究领域(Research Area)

PPAR;Pathways in cancer;Bladder cancer;


图片 (Image Data)

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838




产品货号: APRab09008





Western blot analysis of lysates from A549 cells, treated with etoposide 25uM 24h, using MMP1 (Cleaved-Phe100) Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using Cleaved-MMP-1 22k (F100) Polyclonal Antibody

## 注意事项 (Note)

For research use only.

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838