

产品名称: Cleaved-Notch 2 (D1733) Rabbit Polyclonal Antibody
产品货号: APRab09020



产品概述 (Summary)

产品名称 (Production Name)	Cleaved-Notch 2 (D1733) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	NOTCH2
别名 (Alternative Names)	NOTCH2; Neurogenic locus notch homolog protein 2; Notch 2; hN2
基因 ID (Gene ID)	4853.0
蛋白 ID (SwissProt ID)	Q04721.The antiserum was produced against synthesized peptide derived from human Notch 2. AA range:1684-1733

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	110(cleaved)kDa

产品名称: Cleaved-Notch 2 (D1733) Rabbit Polyclonal Antibody
产品货号: APRab09020



研究背景 (Background)


notch 2(NOTCH2) Homo sapiens This gene encodes a member of the Notch family. Members of this Type 1 transmembrane protein family share structural characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats, and an intracellular domain consisting of multiple, different domain types. Notch family members play a role in a variety of developmental processes by controlling cell fate decisions. The Notch signaling network is an evolutionarily conserved intercellular signaling pathway which regulates interactions between physically adjacent cells. In Drosophila, notch interaction with its cell-bound ligands (delta, serrate) establishes an intercellular signaling pathway that plays a key role in development. Homologues of the notch-ligands have also been identified in human, but precise interactions between these ligands and the human notch homologues remain to be determined. This protein is cleaved. Defects in NOTCH2 are the cause of Alagille syndrome type 2 (ALGS2) [MIM:610205]. Alagille syndrome is an autosomal dominant multisystem disorder defined clinically by hepatic bile duct paucity and cholestasis in association with cardiac, skeletal, and ophthalmologic manifestations. There are characteristic facial features and less frequent clinical involvement of the renal and vascular systems. Functions as a receptor for membrane-bound ligands Jagged1, Jagged2 and Delta1 to regulate cell-fate determination. Upon ligand activation through the released notch intracellular domain (NICD) it forms a transcriptional activator complex with RBP-J kappa and activates genes of the enhancer of split locus. Affects the implementation of differentiation, proliferation and apoptotic programs. PTM: Phosphorylated. PTM: Synthesized in the endoplasmic reticulum as an inactive form which is proteolytically cleaved by a furin-like convertase in the trans-Golgi network before it reaches the plasma membrane to yield an active, ligand-accessible form. Cleavage results in a C-terminal fragment N(TM) and a N-terminal fragment N(EC). Following ligand binding, it is cleaved by TNF-alpha converting enzyme (TACE) to yield a membrane-associated intermediate fragment called notch extracellular truncation (NEXT). This fragment is then cleaved by presenilin dependent gamma-secretase to release a notch-derived peptide containing the intracellular domain (NICD) from the membrane. Similarity: Belongs to the NOTCH family. Similarity: Contains 3 LNR (Lin/Notch) repeats. Similarity: Contains 35 EGF-like domains. Similarity: Contains 6 ANK repeats. Subcellular location: Following proteolytical processing NICD is translocated to the nucleus. Subunit: Heterodimer of a C-terminal fragment N(TM) and an N-terminal fragment N(EC) which are probably linked by disulfide bonds (By similarity). Interacts with MAML1, MAML2 and MAML3 which act as transcriptional coactivators for NOTCH2. Tissue specificity: Expressed in the brain, heart, kidney, lung, skeletal muscle and liver.

研究领域 (Research Area)


Dorso-ventral axis formation; Notch;

图片 (Image Data)


产品名称: Cleaved-Notch 2 (D1733) Rabbit Polyclonal Antibody
产品货号: APRab09020



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using Notch 2 (Cleaved-Asp1733) Antibody.
The picture on the right is blocked with the synthesized peptide.




Western blot analysis of lysates from 293 cells, treated with TNF- α 20ng/ml 30', using Notch 2 (Cleaved-Asp1733) Antibody. The lane on the right is blocked with the synthesized peptide.




Immunofluorescence analysis of rat-lung tissue. 1, Cleaved-Notch 2 (D1733) Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B


产品名称: Cleaved-Notch 2 (D1733) Rabbit Polyclonal Antibody
产品货号: AP Rab09020



Immunofluorescence analysis of rat-lung tissue. 1, Cleaved-Notch 2 (D1733) Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture C: merge of A+B




Immunofluorescence analysis of rat-kidney tissue. 1, Cleaved-Notch 2 (D1733) Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture C: merge of A+B




Immunofluorescence analysis of rat-kidney tissue. 1, Cleaved-Notch 2 (D1733) Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture C: merge of A+B


产品名称: Cleaved-Notch 2 (D1733) Rabbit Polyclonal Antibody
产品货号: APRab09020



Immunohistochemical analysis of paraffin-embedded Human-liver tissue. 1,Cleaved-Notch 2 (D1733) Polyclonal Antibody was diluted at 1:200 (4°C,overnight) . 2, Sodium citrate pH 6.0 was used for antibody retrieval (>98°C,20min) . 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) . Negative control was used by secondary antibody only.



Immunohistochemical analysis of paraffin-embedded Rat-lung tissue. 1,Cleaved-Notch 2 (D1733) Polyclonal Antibody was diluted at 1:200 (4°C,overnight) . 2, Sodium citrate pH 6.0 was used for antibody retrieval (>98°C,20min) . 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) . Negative control was used by secondary antibody only.



Immunohistochemical analysis of paraffin-embedded Rat-spleen tissue. 1,Cleaved-Notch 2 (D1733) Polyclonal Antibody was diluted at 1:200 (4°C,overnight) . 2, Sodium citrate pH 6.0 was used for antibody retrieval (>98°C,20min) . 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) . Negative control was used by secondary antibody only.

产品名称: Cleaved-Notch 2 (D1733) Rabbit Polyclonal
Antibody
产品货号: APRab09020



注意事项 (Note)

For research use only .