产品名称: CMTA2 Rabbit Polyclonal Antibody

产品货号: APRab09083



产品概述 (Summary)

产品名称 (Production Name) CMTA2 Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit 应用 (Application) WB,ELISA

种属反应性 (Reactivity) Human, Mouse

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

储存溶液 (Buffer) Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) CAMTA2 KIAA0909

别名 (Alternative Names)

基因 ID (Gene ID) 23125.0

蛋白 ID (SwissProt ID) O94983.Synthesized peptide derived from part region of human protein

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,ELISA 1:5000-1:20000

蛋白分子量 (Molecular Weight) 132kDa

研究背景 (Background)

The protein encoded by this gene is a member of the calmodulin-binding transcription activator protein family. Members of this family share a common domain structure that consists of a transcription activation domain, a DNA-binding domain,

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: CMTA2 Rabbit Polyclonal Antibody

产品货号: APRab09083



and a calmodulin-binding domain. The encoded protein may be a transcriptional coactivator of genes involved in cardiac growth. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Jan 2010],function:Transcription activator. May act as tumor suppressor.,similarity:Belongs to the CAMTA family.,similarity:Contains 1 CG-1 DNA-binding domain.,similarity:Contains 1 IPT/TIG domain.,similarity:Contains 2 IQ domains.,similarity:Contains 3 ANK repeats.,subunit:May interact with calmodulin .,tissue specificity:Detected in brain. Expressed at constant levels throughout the cell cycle in neuroblastoma cell lines.,

研究领域 (Research Area)

图片 (Image Data)



Western Blot analysis of HEK293 lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838