

产品名称: CNTP4 Rabbit Polyclonal Antibody
产品货号: APRab09141



产品概述 (Summary)

产品名称 (Production Name)	CNTP4 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CNTNAP4 CASPR4 KIAA1763
别名 (Alternative Names)	
基因 ID (Gene ID)	85445.0
蛋白 ID (SwissProt ID)	Q9C0A0.Synthesized peptide derived from part region of human protein

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	143kDa

研究背景 (Background)

This gene encodes a member of the neurexin protein family. Members of this family function in the vertebrate nervous system as cell adhesion molecules and receptors. This protein contains epidermal growth factor repeats and laminin G


产品名称: CNTP4 Rabbit Polyclonal Antibody
产品货号: APRab09141



domains. In addition, it includes an F5/8 type C domain, discoidin/neuropilin- and fibrinogen-like domains, and thrombospondin N-terminal-like domains. This protein may also play a role in proper neurotransmission in the dopaminergic and GABAergic systems and mutations in this gene may be associated with certain psychiatric illnesses. A polymorphism in an intron of this gene may be associated with longevity. [provided by RefSeq, Apr 2016],similarity:Belongs to the neurexin family.,similarity:Contains 1 F5/8 type C domain.,similarity:Contains 1 fibrinogen C-terminal domain.,similarity:Contains 2 EGF-like domains.,similarity:Contains 4 laminin G-like domains.,

研究领域 (Research Area)

图片 (Image Data)



Western Blot analysis of HEK293 lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000

注意事项 (Note)

For research use only .