

产品名称: Cortactin Rabbit Polyclonal Antibody
产品货号: APRab09258



产品概述 (Summary)

产品名称 (Production Name)	Cortactin Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat,Monkey

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CTTN
别名 (Alternative Names)	CTTN; EMS1; Src substrate cortactin; Amplexin; Oncogene EMS1
基因 ID (Gene ID)	13043.0
蛋白 ID (SwissProt ID)	Q14247.The antiserum was produced against synthesized peptide derived from mouse Cortactin. AA range:441-490

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	80kDa

研究背景 (Background)

产品名称: Cortactin Rabbit Polyclonal Antibody
产品货号: AP Rab09258




cortactin(CTTN) Homo sapiens This gene is overexpressed in breast cancer and squamous cell carcinomas of the head and neck. The encoded protein is localized in the cytoplasm and in areas of the cell-substratum contacts. This gene has two roles: (1) regulating the interactions between components of adherens-type junctions and (2) organizing the cytoskeleton and cell adhesion structures of epithelia and carcinoma cells. During apoptosis, the encoded protein is degraded in a caspase-dependent manner. The aberrant regulation of this gene contributes to tumor cell invasion and metastasis. Three splice variants that encode different isoforms have been identified for this gene. [provided by RefSeq, May 2010],function:May contribute to the organization of cell structure. The SH3 motif may function as a binding region to cytoskeleton. Tyrosine phosphorylation in transformed cells may contribute to cellular growth regulation and transformation.,online information:Cortactin entry,similarity:Contains 1 SH3 domain.,similarity:Contains 7 cortactin repeats.,subcellular location:Associated with membrane ruffles and lamellipodia.,subunit:Interacts with SHANK2 and SHANK3 via its SH2 domain. Also interacts with FGD1 (By similarity). Interacts with PLXDC2.,


研究领域 (Research Area)

Tight junction;Pathogenic Escherichia coli infection;

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using Cortactin Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from LOVO cells, using Cortactin Antibody. The lane on the right is blocked with the

产品名称: Cortactin Rabbit Polyclonal Antibody
产品货号: APRab09258



synthesized peptide.

注意事项 (Note)

For research use only .