

产品名称: COX82 Rabbit Polyclonal Antibody
产品货号: APRab09283



产品概述 (Summary)

产品名称 (Production Name)	COX82 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	COX8A
别名 (Alternative Names)	Cytochrome c oxidase subunit 8A, mitochondrial (Cytochrome c oxidase polypeptide VIII-liver/heart;Cytochrome c oxidase subunit 8-2)
基因 ID (Gene ID)	1351.0
蛋白 ID (SwissProt ID)	P10176.Synthesized peptide derived from human COX82 AA range: 10-90

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:50-1:200,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	

研究背景 (Background)


产品名称: COX82 Rabbit Polyclonal Antibody
产品货号: APRab09283



The protein encoded by this gene is the terminal enzyme of the respiratory chain, coupling the transfer of electrons from cytochrome c to molecular oxygen, with the concomitant production of a proton electrochemical gradient across the inner mitochondrial membrane. In addition to 3 mitochondrially encoded subunits, which perform the catalytic function, the eukaryotic enzyme contains nuclear-encoded smaller subunits, ranging in number from 4 in some organisms to 10 in mammals. It has been proposed that nuclear-encoded subunits may be involved in the modulation of the catalytic function. This gene encodes one of the nuclear-encoded subunits. [provided by RefSeq, Jul 2008],function:This protein is one of the nuclear-coded polypeptide chains of cytochrome c oxidase, the terminal oxidase in mitochondrial electron transport.,similarity:Belongs to the cytochrome c oxidase VIII family,.

研究领域 (Research Area)

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200 (4° overnight) . 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .

注意事项 (Note)

For research use only .