产品名称: CRLF3 Rabbit Polyclonal Antibody

产品货号: APRab09408



产品概述 (Summary)

产品名称 (Production Name) CRLF3 Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host)Rabbit应用 (Application)WB,ELISA种属反应性 (Reactivity)Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

储存溶液 (Buffer) Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) CRLF3 CREME9 CYTOR4 P48

别名 (Alternative Names)

基因 ID (Gene ID) 51379.0

Q8IUI8.Synthesized peptide derived from human protein . at AA range: 270-蛋白ID (SwissProt ID)

350

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,ELISA 1:5000-1:20000

蛋白分子量 (Molecular Weight) 48kDa

研究背景 (Background)

This gene encodes a cytokine receptor-like factor that may negatively regulate cell cycle progression at the G0/G1 phase.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: CRLF3 Rabbit Polyclonal Antibody


产品货号: APRab09408



Studies of the related rat protein suggest that it may regulate neuronal morphology and synaptic vesicle biogenesis. This gene is one of several genes located in the neurofibromatosis type I tumor suppressor region on the q arm of chromosome 17, a region that is subject to microdeletions, duplications, chromosomal breaks and rearrangements. Alternative splicing of this gene results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 2 and 5. [provided by RefSeq, Aug 2012], similarity: Belongs to the cytokine receptor-like factor 3 family., similarity: Contains 1 fibronectin type-III domain., tissue specificity: Expressed in skin and squamous cell carcinoma (SCC) and in lesion actinic keratosis (AK).,

研究领域 (Research Area)

图片 (Image Data)



Western Blot analysis of Hela lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838