

产品名称: Crystallin- α B Rabbit Polyclonal Antibody
产品货号: APRab09439



产品概述 (Summary)

产品名称 (Production Name)	Crystallin- α B Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CRYAB CRYAB; CRYA2; Alpha-crystallin B chain; Alpha(B)-crystallin; Heat shock protein beta-5; HspB5; Renal carcinoma antigen NY-REN-27; Rosenthal fiber component
别名 (Alternative Names)	
基因 ID (Gene ID)	1410.0
蛋白 ID (SwissProt ID)	P02511.The antiserum was produced against synthesized peptide derived from human CRYAB. AA range:10-59

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:10000
蛋白分子量 (Molecular Weight)	24kDa

产品名称: Crystallin- α B Rabbit Polyclonal Antibody
产品货号: APRab09439



研究背景 (Background)


Mammalian lens crystallins are divided into alpha, beta, and gamma families. Alpha crystallins are composed of two gene products: alpha-A and alpha-B, for acidic and basic, respectively. Alpha crystallins can be induced by heat shock and are members of the small heat shock protein (HSP20) family. They act as molecular chaperones although they do not renature proteins and release them in the fashion of a true chaperone; instead they hold them in large soluble aggregates. Post-translational modifications decrease the ability to chaperone. These heterogeneous aggregates consist of 30-40 subunits; the alpha-A and alpha-B subunits have a 3:1 ratio, respectively. Two additional functions of alpha crystallins are an autokinase activity and participation in the intracellular architecture. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. Crystallins do not turn over as the lens ages, providing ample opportunity for post-translational modifications or oxidations. These modifications may change crystallin solubility properties and favor senile cataract. Defects in CRYAB are the cause of alpha-B crystallinopathy [MIM:608810]. Alpha-B crystallinopathy is an autosomal dominant form of desmin-related myopathy (DRM) that results in weakness of the proximal and distal limb muscle (including neck, velopharynx, and trunk muscles), signs of cardiomyopathy and cataract. Patients with progressive myopathy characterized by myofibrillar degeneration that commences at the Z-disk, have been described. Mutations truncate the essential C-terminal domain of the protein required for the chaperone function. Seen as Rosenthal fiber protein in the brain tissue of patients with Alexander disease. May contribute to the transparency and refractive index of the lens. mass spectrometry: PubMed:10930324, mass spectrometry: PubMed:8175657, mass spectrometry: With 1 phosphate group PubMed:10930324, mass spectrometry: With 1 phosphate group PubMed:8175657, mass spectrometry: With 2 phosphate groups PubMed:8175657, similarity: Belongs to the small heat shock protein (HSP20) family. subunit: Aggregates with homologous proteins, including CRYAA and the small heat shock protein HSPB1, to form large heteromeric complexes. Interacts with HSPBAP1 and TTN/titin. tissue specificity: Lens as well as other tissues.

研究领域 (Research Area)


Signal Transduction

图片 (Image Data)


产品名称: Crystallin- α B Rabbit Polyclonal Antibody
产品货号: APRab09439



Immunohistochemistry analysis of paraffin-embedded human skeletal muscle tissue, using CRYAB Antibody. The picture on the right is blocked with the synthesized peptide.




Western blot analysis of lysates from 293 cells, using CRYAB Antibody. The lane on the right is blocked with the synthesized peptide.




Western Blot analysis of various cells using Crystallin- α B Polyclonal Antibody


产品名称: Crystallin- α B Rabbit Polyclonal Antibody
产品货号: APRab09439



Western Blot analysis of 293 cells using Crystallin- α B Polyclonal Antibody




Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:400 (4°, overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 30min) .



Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:400 (4°, overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 30min) .

产品名称: Crystallin- α B Rabbit Polyclonal Antibody
产品货号: APRab09439



Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:400 (4°, overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 30min) .

注意事项 (Note)

For research use only .