

产品名称: CUL-2 Rabbit Polyclonal Antibody
产品货号: APRab09532



产品概述 (Summary)

产品名称 (Production Name)	CUL-2 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CUL2
别名 (Alternative Names)	CUL2; Cullin-2; CUL-2
基因 ID (Gene ID)	8453.0
蛋白 ID (SwissProt ID)	Q13617.The antiserum was produced against synthesized peptide derived from human Cullin 2. AA range:696-745

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	87kDa

研究背景 (Background)

产品名称: CUL-2 Rabbit Polyclonal Antibody
产品货号: AP Rab09532



function:Core component of multiple cullin-RING-based ECS (ElonginB/C-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complexes, which mediate the ubiquitination of target proteins. May serve as a rigid scaffold in the complex and may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme. The E3 ubiquitin-protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1 (By similarity). The functional specificity of the ECS complex depends on the substrate recognition component. ECS(VHL) mediates the ubiquitination of hypoxia-inducible factor (HIF).,pathway:Protein modification; protein ubiquitination.,PTM:CBC(VHL) complex formation seems to promote neddylation. Deneddylation via its interaction with the COP9 signalosome (CSN) complex.,similarity:Belongs to the cullin family.,subunit:Component of multiple ECS (Elongin BC-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complexes formed of CUL2, Elongin BC (TCEB1 and TCEB2), RBX1 and a variable substrate-specific adapter. Component of the ECS(VHL) or CBC(VHL) complex containing VHL. Component of the ECS(MED8) complex with the probable substrate recognition component MED8 (By similarity). Component of the ECS(PPIL5) complex with the probable substrate recognition component PPIL5. Component of a probable ECS E3 ubiquitin-protein ligase complex containing CUL2, RBX1, TCEB1, TCEB2 and FEM1B. Part of an E3 ubiquitin-protein ligase complex including ZYG11B, CUL2 and Elongin BC. Part of an E3 ubiquitin-protein ligase complex including ZYG11BL, CUL2 and Elongin BC.Interacts with RBX1, RNF7, FEM1B and TIP120A/CAND1. Interacts with COPS2, and MED8 (By similarity). Interacts with human respiratory syncytial virus (HRSV) protein NS1.,function:Core component of multiple cullin-RING-based ECS (ElonginB/C-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complexes, which mediate the ubiquitination of target proteins. May serve as a rigid scaffold in the complex and may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme. The E3 ubiquitin-protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1 (By similarity). The functional specificity of the ECS complex depends on the substrate recognition component. ECS(VHL) mediates the ubiquitination of hypoxia-inducible factor (HIF).,pathway:Protein modification; protein ubiquitination.,PTM:CBC(VHL) complex formation seems to promote neddylation. Deneddylation via its interaction with the COP9 signalosome (CSN) complex.,similarity:Belongs to the cullin family.,subunit:Component of multiple ECS (Elongin BC-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complexes formed of CUL2, Elongin BC (TCEB1 and TCEB2), RBX1 and a variable substrate-specific adapter. Component of the ECS(VHL) or CBC(VHL) complex containing VHL. Component of the ECS(MED8) complex with the probable substrate recognition component MED8 (By similarity). Component of the ECS(PPIL5) complex with the probable substrate recognition component PPIL5. Component of a probable ECS E3 ubiquitin-protein ligase complex containing CUL2, RBX1, TCEB1, TCEB2 and FEM1B. Part of an E3 ubiquitin-protein ligase complex including ZYG11B, CUL2 and Elongin BC. Part of an E3 ubiquitin-protein ligase complex including ZYG11BL, CUL2 and Elongin BC.Interacts with RBX1, RNF7, FEM1B and TIP120A/CAND1. Interacts with COPS2, and MED8 (By similarity). Interacts with human respiratory syncytial virus (HRSV) protein NS1.,


研究领域 (Research Area)

Ubiquitin mediated proteolysis;Pathways in cancer;Renal cell carcinoma;


产品名称: CUL-2 Rabbit Polyclonal Antibody
产品货号: APRab09532




图片 (Image Data)



Immunofluorescence analysis of NIH/3T3 cells, using Cullin 2 Antibody. The picture on the right is blocked with the synthesized peptide.




Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using Cullin 2 Antibody. The picture on the right is blocked with the synthesized peptide.




Western blot analysis of lysates from LOVO cells, using Cullin 2 Antibody. The lane on the right is blocked with the synthesized peptide.

产品名称: CUL-2 Rabbit Polyclonal Antibody
产品货号: APRab09532



Western Blot analysis of various cells using CUL-2 Polyclonal Antibody diluted at 1: 2000



Western Blot analysis of Jurkat cells using CUL-2 Polyclonal Antibody diluted at 1: 2000

注意事项 (Note)

For research use only .