

产品名称: Cyclin A1 Rabbit Polyclonal Antibody
产品货号: APRab09578



产品概述 (Summary)

产品名称 (Production Name)	Cyclin A1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CCNA1
别名 (Alternative Names)	CCNA1; Cyclin-A1
基因 ID (Gene ID)	8900.0
蛋白 ID (SwissProt ID)	P78396.The antiserum was produced against synthesized peptide derived from human Cyclin A1. AA range:411-460

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	52kDa

研究背景 (Background)

产品名称: Cyclin A1 Rabbit Polyclonal Antibody
产品货号: AP Rab09578




The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. The cyclin encoded by this gene was shown to be expressed in testis and brain, as well as in several leukemic cell lines, and is thought to primarily function in the control of the germline meiotic cell cycle. This cyclin binds both CDK2 and CDC2 kinases, which give two distinct kinase activities, one appearing in S phase, the other in G2, and thus regulate separate functions in cell cycle. This cyclin was found to bind to important cell cycle regulators, such as Rb family proteins, transcription factor E2F-1, and the p21 family proteins. Multidevelopmental stage: Expression increases in early G1 phase and reaches highest levels during the S and G2/M phases. function: May be involved in the control of the cell cycle at the G1/S (start) and G2/M (mitosis) transitions. May primarily function in the control of the germline meiotic cell cycle and additionally in the control of mitotic cell cycle in some somatic cells. similarity: Belongs to the cyclin family. similarity: Belongs to the cyclin family. Cyclin AB subfamily. subunit: Interacts with the CDK2 and the CDC2 protein kinases to form a serine/threonine kinase holoenzyme complex. The cyclin subunit imparts substrate specificity to the complex. Does not bind CDK4 and CDK5 (in vitro). The cyclin A1-CDK2 complex interacts with transcription factor E2F-1 and RB proteins. Found in a complex with CDK2, CABLES1 and CCNE1 (By similarity). Interacts with INCA1 and KLHDC9. tissue specificity: Very high levels in testis and very low levels in brain. Also found in myeloid Leukemia cell lines.

研究领域 (Research Area)


AMPK

图片 (Image Data)




Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using Cyclin A1 Antibody. The picture on the right is blocked with the synthesized peptide.


产品名称: Cyclin A1 Rabbit Polyclonal Antibody
产品货号: APRab09578



Western blot analysis of lysates from SKOV3 cells, using Cyclin A1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using Cyclin A1 Polyclonal Antibody diluted at 1: 2000



Western Blot analysis of HeLa cells using Cyclin A1 Polyclonal Antibody diluted at 1: 2000

注意事项 (Note)

For research use only .