

产品名称: CYP2B6 Rabbit Polyclonal Antibody
产品货号: APRab09650



产品概述 (Summary)

产品名称 (Production Name)	CYP2B6 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CYP2B6
别名 (Alternative Names)	CYP2B6; Cytochrome P450 2B6; 1; 4-cineole 2-exo-monooxygenase; CYP11B6; Cytochrome P450 IIB1
基因 ID (Gene ID)	1565.0
蛋白 ID (SwissProt ID)	P20813.Synthesized peptide derived from the Internal region of human CYP2B6.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	55kDa

产品名称: CYP2B6 Rabbit Polyclonal Antibody
产品货号: AP Rab09650




研究背景 (Background)

This gene, CYP2B6, encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by phenobarbital. The enzyme is known to metabolize some xenobiotics, such as the anti-cancer drugs cyclophosphamide and ifosfamide. Transcript variants for this gene have been described; however, it has not been resolved whether these transcripts are in fact produced by this gene or by a closely related pseudogene, CYP2B7. Both the gene and the pseudogene are located in the middle of a CYP2A pseudogene found in a large cluster of cytochrome P450 genes from the CYP2A, CYP2B and CYP2F subfamilies on chromosome 19q. [provided by RefSeq, Jul 2008], catalytic activity: $\text{RH} + \text{reduced flavoprotein} + \text{O}_2 = \text{ROH} + \text{oxidized flavoprotein} + \text{H}_2\text{O}$, cofactor: Heme group, function: Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver microsomes, this enzyme is involved in an NADPH-dependent electron transport pathway. It oxidizes a variety of structurally unrelated compounds, including steroids, fatty acids, and xenobiotics, induction: By phenobarbital, online information: CYP2B6 alleles, PTM: Phosphorylation is accompanied by a decrease in enzyme activity, similarity: Belongs to the cytochrome P450 family, tissue specificity: Expressed in liver, lung and heart right ventricle,

研究领域 (Research Area)

Arachidonic acid metabolism; Retinol metabolism; Metabolism of xenobiotics by cytochrome P450; Drug metabolism;

图片 (Image Data)



Western Blot analysis of 293 cells using CYP2B6 Polyclonal Antibody

注意事项 (Note)

For research use only .