

产品名称: CYP2C8 Rabbit Polyclonal Antibody
产品货号: APRab09654



产品概述 (Summary)

产品名称 (Production Name)	CYP2C8 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CYP2C8 CYP2C8; Cytochrome P450 2C8; CYP11C8; Cytochrome P450 11C2; Cytochrome
别名 (Alternative Names)	P450 MP-12; Cytochrome P450 MP-20; Cytochrome P450 form 1; S-mephenytoin 4-hydroxylase
基因 ID (Gene ID)	1558.0
蛋白 ID (SwissProt ID)	P10632.Synthesized peptide derived from the Internal region of human CYP2C8.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	55kDa

产品名称: CYP2C8 Rabbit Polyclonal Antibody
产品货号: APRab09654




研究背景 (Background)

This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by phenobarbital. The enzyme is known to metabolize many xenobiotics, including the anticonvulsive drug mephenytoin, benzo(a)pyrene, 7-ethoxycoumarin, and the anti-cancer drug taxol. This gene is located within a cluster of cytochrome P450 genes on chromosome 10q24. Several transcript variants encoding a few different isoforms have been found for this gene. [provided by RefSeq, Nov 2010], catalytic activity: $\text{RH} + \text{reduced flavoprotein} + \text{O}_2 = \text{ROH} + \text{oxidized flavoprotein} + \text{H}_2\text{O}$, caution: Alternative splicing has been shown to occur but the shorter forms are believed to be non-functional., cofactor: Heme group., function: Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver microsomes, this enzyme is involved in an NADPH-dependent electron transport pathway. It oxidizes a variety of structurally unrelated compounds, including steroids, fatty acids, and xenobiotics. In the epoxidation of arachidonic acid it generates only 14,15- and 11,12-cis-epoxyeicosatrienoic acids. It is the principal enzyme responsible for the metabolism the anti-cancer drug paclitaxel (taxol)., induction: By phenobarbital., online information: CYP2C8 alleles, similarity: Belongs to the cytochrome P450 family.,

研究领域 (Research Area)

Arachidonic acid metabolism; Linoleic acid metabolism; Retinol metabolism; Metabolism of xenobiotics by cytochrome P450; Drug metabolism;

图片 (Image Data)



Western Blot analysis of 293 cells using CYP2C8 Polyclonal Antibody diluted at 1: 1000

注意事项 (Note)

For research use only .