

产品名称: DAP-5 Rabbit Polyclonal Antibody
产品货号: APRab09785



产品概述 (Summary)

产品名称 (Production Name)	DAP-5 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	EIF4G2
别名 (Alternative Names)	EIF4G2; DAP5; OK/SW-cl.75; Eukaryotic translation initiation factor 4 gamma 2; eIF-4-gamma 2; eIF-4G 2; eIF4G 2; Death-associated protein 5; DAP-5; p97
基因 ID (Gene ID)	1982.0
蛋白 ID (SwissProt ID)	P78344.The antiserum was produced against synthesized peptide derived from human EIF4G2. AA range:41-90

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	90kDa

产品名称: DAP-5 Rabbit Polyclonal Antibody
产品货号: APRab09785




研究背景 (Background)

Translation initiation is mediated by specific recognition of the cap structure by eukaryotic translation initiation factor 4F (eIF4F), which is a cap binding protein complex that consists of three subunits: eIF4A, eIF4E and eIF4G. The protein encoded by this gene shares similarity with the C-terminal region of eIF4G that contains the binding sites for eIF4A and eIF3; eIF4G, in addition, contains a binding site for eIF4E at the N-terminus. Unlike eIF4G, which supports cap-dependent and independent translation, this gene product functions as a general repressor of translation by forming translationally inactive complexes. In vitro and in vivo studies indicate that translation of this mRNA initiates exclusively at a non-AUG (GUG) codon. Alternatively spliced transcript variants encoding different isoforms of this gene have been described. [provided by RefSeq, Jul 2008],function:Appears to play a role in the switch from cap-dependent to IRES-mediated translation during mitosis, apoptosis and viral infection. Cleaved by some caspases and viral proteases.,miscellaneous:This gene has been shown to be extensively edited in the liver of APOBEC1 transgenic animal model. Its aberrant editing could contribute to the potent oncogenesis induced by overexpression of APOBEC1. The aberrant edited sequence, called NAT1, is likely to be a fundamental translational repressor.,PTM:Phosphorylation; hyperphosphorylated during mitosis.,similarity:Belongs to the eIF4G family.,similarity:Contains 1 MI domain.,similarity:Contains 1 MIF4G domain.,similarity:Contains 1 W2 domain.,subunit:Interacts with the serine/threonine protein kinases MKNK1 and MKNK2. Binds EIF4A and EIF3. Interacts with MIF4GD.,tissue specificity:Ubiquitously expressed in all adult tissues examined, with high levels in skeletal muscle and heart. Also expressed in fetal brain, lung, liver and kidney.,

研究领域 (Research Area)


Viral myocarditis;

图片 (Image Data)




Western blot analysis of lysates from A549 cells, using EIF4G2 Antibody. The lane on the right is blocked with the synthesized peptide.

产品名称: DAP-5 Rabbit Polyclonal Antibody
产品货号: APRab09785



Western Blot analysis of various cells using DAP-5 Polyclonal Antibody



Western Blot analysis of A549 cells using DAP-5 Polyclonal Antibody

注意事项 (Note)

For research use only .