产品名称: DCAMKL2 Rabbit Polyclonal Antibody

产品货号: APRab09829



## 产品概述 (Summary)

产品名称 (Production Name) DCAMKL2 Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application)WB,ICC/IF,ELISA种属反应性 (Reactivity)Human,Mouse

### 产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

# 免疫原信息 (Immunogen)

基因名 (Gene Name) DCLK2

DCLK2; DCAMKL2; DCDC3B; DCK2; Serine/threonine-protein kinase DCLK2;

CaMK-like CREB regulatory kinase 2; CL2; CLICK-II; CLICK2; Doublecortin

别名 (Alternative Names)

domain-containing protein 3B; Doublecortin-like and CAM kinase-like 2;

domain-containing protein 3b, Doublecortin-like and CAW kinase-like 2

Doublecortin-like kinase 2

基因 ID (Gene ID) 166614.0

Q8N568.The antiserum was produced against synthesized peptide derived 蛋白ID (SwissProt ID)

from human DCLK2. AA range:1-50

# 产品应用 (Application)

**稀释比 (Dilution Ratio)** WB 1:500-1:2000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: DCAMKL2 Rabbit Polyclonal Antibody

产品货号: APRab09829




**蛋白分子量 (Molecular Weight)** 83kDa

#### 研究背景 (Background)

This gene encodes a member of the protein kinase superfamily and the doublecortin family. The protein encoded by this gene contains two N-terminal doublecortin domains, which bind microtubules and regulate microtubule polymerization, a C-terminal serine/threonine protein kinase domain, which shows substantial homology to Ca2+/calmodulin-dependent protein kinase, and a serine/proline-rich domain in between the doublecortin and the protein kinase domains, which mediates multiple protein-protein interactions. The microtubule-polymerizing activity of the encoded protein is independent of its protein kinase activity. Mouse studies show that the DCX gene, another family member, and this gene share function in the establishment of hippocampal organization and that their absence results in a severe epileptic phenotype and lethality, as described in human patients with lissencephaly. Multiple alternacatalytic activity:ATP + a protein = ADP + a phosphoprotein, similarity:Belongs to the protein kinase superfamily, similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. CaMK subfamily, similarity:Contains 1 protein kinase domain, similarity:Contains 2 doublecortin domains,

## 研究领域 (Research Area)

#### 图片 (Image Data)




Western blot analysis of lysates from HepG2 cells, using DCLK2 Antibody. The lane on the right is blocked with the synthesized peptide.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品名称: DCAMKL2 Rabbit Polyclonal Antibody

产品货号: APRab09829





Western Blot analysis of various cells using DCAMKL2 Polyclonal Antibody

## 注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838