

产品名称: DCC Rabbit Polyclonal Antibody
产品货号: APRab09832



产品概述 (Summary)

产品名称 (Production Name)	DCC Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	DCC DCC; IGDCC1; Netrin receptor DCC; Colorectal cancer suppressor;
别名 (Alternative Names)	Immunoglobulin superfamily DCC subclass member 1; Tumor suppressor protein DCC
基因 ID (Gene ID)	1630.0
蛋白 ID (SwissProt ID)	P43146.The antiserum was produced against synthesized peptide derived from human DCC. AA range:441-490

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	160kDa

产品名称: DCC Rabbit Polyclonal Antibody
产品货号: AP Rab09832




研究背景 (Background)

This gene encodes a netrin 1 receptor. The transmembrane protein is a member of the immunoglobulin superfamily of cell adhesion molecules, and mediates axon guidance of neuronal growth cones towards sources of netrin 1 ligand. The cytoplasmic tail interacts with the tyrosine kinases Src and focal adhesion kinase (FAK, also known as PTK2) to mediate axon attraction. The protein partially localizes to lipid rafts, and induces apoptosis in the absence of ligand. The protein functions as a tumor suppressor, and is frequently mutated or downregulated in colorectal cancer and esophageal carcinoma. [provided by RefSeq, Oct 2009],disease:Colorectal tumors that lost their capacity to differentiate into mucus producing cells uniformly lack DCC expression. Inactivation of DCC due to allelic deletion and/or point mutations may cause both lymphatic and hematogenous metastasis of esophageal squamous cell carcinomas.,function:Receptor for netrin required for axon guidance. Mediates axon attraction of neuronal growth cones in the developing nervous system upon ligand binding. Its association with UNC5 proteins may trigger signaling for axon repulsion. It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand. Implicated as a tumor suppressor gene.,PTM:Ubiquitinated; mediated by SIAH1 or SIAH2 and leading to its subsequent proteasomal degradation.,similarity:Belongs to the immunoglobulin superfamily. DCC family.,similarity:Contains 4 Ig-like C2-type (immunoglobulin-like) domains.,similarity:Contains 6 fibronectin type-III domains.,subunit:Interacts with the cytoplasmic part of UNC5A, UNC5B, UNC5C and probably UNC5D.,tissue specificity:Found in axons of the central and peripheral nervous system and in differentiated cell types of the intestine.,

研究领域 (Research Area)


Axon guidance;Pathways in cancer;Colorectal cancer;

图片 (Image Data)



Immunofluorescence analysis of LOVO cells, using DCC Antibody. The picture on the right is blocked with the synthesized peptide.

产品名称: DCC Rabbit Polyclonal Antibody
产品货号: APRab09832



Western Blot analysis of 293 cells using DCC Polyclonal Antibody diluted at 1: 500

注意事项 (Note)

For research use only .