

产品名称: DDR1 Rabbit Polyclonal Antibody
产品货号: AP Rab09868



产品概述 (Summary)

产品名称 (Production Name)	DDR1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	DDR1 CAK EDDR1 NEP NTRK4 PTK3A RTK6 TRKE Epithelial discoidin domain-containing receptor 1 (Epithelial discoidin domain receptor 1) (EC 2.7.10.1) (CD167 antigen-like family member A) (Cell adhesion kinase) (Discoidin receptor tyrosine kinase) (HGK2) (Mammary carcinoma kinase 10) (MCK-10) (Protein-tyrosine kinase 3A) (Protein-tyrosine kinase RTK-6) (TRK E) (Tyrosine kinase DDR) (Tyrosine-protein kinase CAK) (CD antigen CD167a)
别名 (Alternative Names)	
基因 ID (Gene ID)	780.0
蛋白 ID (SwissProt ID)	Q08345.Synthesized peptide derived from human DDR1

产品应用 (Application)

产品名称: DDR1 Rabbit Polyclonal Antibody
产品货号: AP Rab09868




稀释比 (Dilution Ratio) WB 1:500-1:2000, ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight) 130kDa

研究背景 (Background)

Receptor tyrosine kinases play a key role in the communication of cells with their microenvironment. These kinases are involved in the regulation of cell growth, differentiation and metabolism. The protein encoded by this gene belongs to a subfamily of tyrosine kinase receptors with homology to Dictyostelium discoideum protein discoidin I in their extracellular domain, and that are activated by various types of collagen. Expression of this protein is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, it has been shown to be significantly overexpressed in several human tumors. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Feb 2011], catalytic activity: ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate., domain: The Gly/Pro-rich domains may be required for an unusual geometry of interaction with ligand or substrates., function: May be involved in cell-cell interactions and recognition., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily., similarity: Contains 1 F5/8 type C domain., similarity: Contains 1 protein kinase domain., tissue specificity: Expressed at low levels in most adult tissues and is highest in the brain and lung. Abundant in breast carcinoma cell lines.,

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of various lysate, antibody was diluted at 1000. Secondary antibody was diluted at 1:20000

注意事项 (Note)

For research use only .