

产品名称: DDR2 Rabbit Polyclonal Antibody
产品货号: APRab09869



产品概述 (Summary)

产品名称 (Production Name)	DDR2 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC/IF, ELISA
种属反应性 (Reactivity)	Human, Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	DDR2 NTRKR3 TKT TYRO10 Discoidin domain-containing receptor 2 (Discoidin domain receptor 2) (EC 2.7.10.1) (CD167 antigen-like family member B) (Discoidin domain-containing receptor tyrosine kinase 2) (Neurotrophic tyrosine kinase, receptor-related 3) (Receptor protein-tyrosine kinase TKT) (Tyrosine-protein kinase TYRO10) (CD antigen CD167b)
别名 (Alternative Names)	
基因 ID (Gene ID)	4921.0
蛋白 ID (SwissProt ID)	Q16832.Synthetic peptide from human protein at AA range: 31-80

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:50-1:200, ICC/IF 1:200-1:500, ELISA 1:10000-1:20000
----------------------	---

产品名称: DDR2 Rabbit Polyclonal Antibody
产品货号: APRab09869




蛋白分子量 (Molecular Weight)

研究背景 (Background)

Receptor tyrosine kinases (RTKs) play a key role in the communication of cells with their microenvironment. These molecules are involved in the regulation of cell growth, differentiation, and metabolism. In several cases the biochemical mechanism by which RTKs transduce signals across the membrane has been shown to be ligand induced receptor oligomerization and subsequent intracellular phosphorylation. This autophosphorylation leads to phosphorylation of cytosolic targets as well as association with other molecules, which are involved in pleiotropic effects of signal transduction. RTKs have a tripartite structure with extracellular, transmembrane, and cytoplasmic regions. This gene encodes a member of a novel subclass of RTKs and contains a distinct extracellular region encompassing a factor VIII-like domain. Alternative splicing in the 5' UTR results in multiple transcatalytic activity: $\text{ATP} + \text{a [protein]-L-tyrosine} = \text{ADP} + \text{a [protein]-L-tyrosine phosphate}$.
function: This tyrosine kinase receptor for fibrillar collagen mediates fibroblast migration and proliferation. Contributes to cutaneous wound healing.
similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily.
similarity: Contains 1 F5/8 type C domain.
similarity: Contains 1 protein kinase domain.
tissue specificity: The major 10 kDa transcript is expressed in high levels in heart and lung, less in brain, placenta, liver, skeletal muscle, pancreas, and kidney.


研究领域 (Research Area)

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human-lung-cancer, antibody was diluted at 1:200

产品名称: DDR2 Rabbit Polyclonal Antibody
产品货号: APRab09869



Immunohistochemical analysis of paraffin-embedded human-lung-cancer, antibody was diluted at 1:200

注意事项 (Note)

For research use only .