

产品名称: EDG-1 Rabbit Polyclonal Antibody
产品货号: APRab10295



产品概述 (Summary)

产品名称 (Production Name)	EDG-1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	S1PR1 S1PR1; CHEDG1; EDG1; Sphingosine 1-phosphate receptor 1; S1P receptor 1;
别名 (Alternative Names)	S1P1; Endothelial differentiation G-protein coupled receptor 1; Sphingosine 1-phosphate receptor Edg-1; S1P receptor Edg-1; CD antigen CD363
基因 ID (Gene ID)	1901.0
蛋白 ID (SwissProt ID)	P21453.The antiserum was produced against synthesized peptide derived from human S1P Receptor EDG1. AA range:206-255

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:10000
蛋白分子量 (Molecular Weight)	50kDa

产品名称: EDG-1 Rabbit Polyclonal Antibody
产品货号: APRab10295




研究背景 (Background)

The protein encoded by this gene is structurally similar to G protein-coupled receptors and is highly expressed in endothelial cells. It binds the ligand sphingosine-1-phosphate with high affinity and high specificity, and suggested to be involved in the processes that regulate the differentiation of endothelial cells. Activation of this receptor induces cell-cell adhesion. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016],function:Receptor for the lysosphingolipid sphingosine 1-phosphate (S1P). S1P is a bioactive lysophospholipid that elicits diverse physiological effect on most types of cells and tissues. This inducible epithelial cell G-protein-coupled receptor may be involved in the processes that regulate the differentiation of endothelial cells. Seems to be coupled to the G(i) subclass of heteromeric G proteins.,induction:By the tumor promoter phorbol 12-myristate 13-acetate (PME) in the presence of cycloheximide.,PTM:S1P-induced endothelial cell migration requires the PKB/AKT1-mediated phosphorylation of the third intracellular loop at the Thr-236 residue.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Endothelial cells, and to a lesser extent, in vascular smooth muscle cells, fibroblasts, melanocytes, and cells of epithelioid origin.,

研究领域 (Research Area)


Neuroactive ligand-receptor interaction;

图片 (Image Data)




Immunofluorescence analysis of A549 cells, using S1P Receptor EDG1 Antibody. The picture on the right is blocked with the synthesized peptide.


产品名称: EDG-1 Rabbit Polyclonal Antibody
产品货号: APRab10295



Western blot analysis of lysates from HT-29, LOVO, and A549 cells, using S1P Receptor EDG1 Antibody. The lane on the right is blocked with the synthesized peptide.




Western blot analysis of the lysates from HUVEC cells using S1P Receptor EDG1 antibody.



Western Blot analysis of various cells using EDG-1 Polyclonal Antibody

产品名称: EDG-1 Rabbit Polyclonal Antibody
产品货号: APRab10295



Western Blot analysis of A549 cells using EDG-1 Polyclonal Antibody

注意事项 (Note)

For research use only .