

**产品名称:** EGF Rabbit Polyclonal Antibody

**产品货号:** APRab10332

### 产品概述 (Summary)

|                   |                                                                                                   |
|-------------------|---------------------------------------------------------------------------------------------------|
| 描述(Description)   | Rabbit polyclonal Antibody                                                                        |
| 宿主(Host)          | Rabbit                                                                                            |
| 应用(Application)   | WB,IHC,ICC/IF,ELISA                                                                               |
| 种属反应(Reactivity)  | Human,Mouse                                                                                       |
| 偶联物(Conjugation)  | Unconjugated                                                                                      |
| 修饰(Modification)  | Unmodified                                                                                        |
| 同种型(Isotype)      | IgG                                                                                               |
| 克隆(Clonality)     | Polyclonal                                                                                        |
| 剂型(Form)          | Liquid                                                                                            |
| 保存条件(Storage)     | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.          |
| 储存溶液(Buffer)      | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| 纯化 (Purification) | Affinity purification                                                                             |

### 产品应用(Application)

|                       |                                                                        |
|-----------------------|------------------------------------------------------------------------|
| 稀释比(Dilution Ratio)   | WB 1:500-1:2000,IHC 1:50-1:200,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000 |
| 分子量(Molecular Weight) | 130kDa                                                                 |

### 抗原信息(Antigen Information)

|                       |                                                                                         |
|-----------------------|-----------------------------------------------------------------------------------------|
| 基因名(Gene Name)        | EGF                                                                                     |
| 别名(Alternative Names) | Pro-epidermal growth factor (EGF) [Cleaved into: Epidermal growth factor (Urogastrone)] |
| 基因 ID(Gene ID)        | 1950.0                                                                                  |
| SwissProt ID          | P01133                                                                                  |
| 免疫原(Immunogen)        | Synthetic peptide from human protein at AA range: 1000-1060                             |

### 研究背景 (Background)

This gene encodes a member of the epidermal growth factor superfamily. The encoded preproprotein is proteolytically processed to generate the 53-amino acid epidermal growth factor peptide. This protein acts a potent mitogenic factor that plays an important role in the growth, proliferation and differentiation of numerous cell types. This protein acts by binding with

high affinity to the cell surface receptor, epidermal growth factor receptor. Defects in this gene are the cause of hypomagnesemia type 4. Dysregulation of this gene has been associated with the growth and progression of certain cancers. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed. [provided by RefSeq, Jan 2016],disease:Defects in EGF are the cause of hypomagnesemia type 4 (HOMG4) [MIM:611718]; also known as renal hypomagnesemia normocalciuric. HOMG4 is a disorder characterized by massive renal hypomagnesemia and normal levels of serum calcium and calcium excretion. Clinical features include seizures, mild-to moderate psychomotor retardation, and brisk tendon reflexes.,function:EGF stimulates the growth of various epidermal and epithelial tissues in vivo and in vitro and of some fibroblasts in cell culture. Magnesiotropic hormone that stimulates magnesium reabsorption in the renal distal convoluted tubule via engagement of EGFR and activation of the magnesium channel TRPM6.,online information:Epidermal growth factor entry,similarity:Contains 9 EGF-like domains.,similarity:Contains 9 LDL-receptor class B repeats.,tissue specificity:Expressed in kidney, salivary gland, cerebrum and prostate.,


### 研究领域 (Research Area)

MAPK\_ERK\_Growth;MAPK\_G\_Protein;ErbB\_HER;Cytokine-cytokine receptor interaction;Endocytosis;Focal adhesion;Gap junction;Regulates Actin and Cytoskeleton;Pathways in cancer;Pancreatic cancer;Endometrial cancer;Glioma;Prostate cancer;Melanoma;Bladder cancer;Non-small cell lung cancer;


### 注意事项 (Note)

For research use only.


### 图片 (Image Data)




Western blot analysis of 293T mouse-kidney lysate, antibody was diluted at 2000. Secondary antibody was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human-kidney-cancer, antibody was diluted at 1:200




Immunohistochemical analysis of paraffin-embedded human-kidney-cancer, antibody was diluted at 1:200




Immunohistochemical analysis of paraffin-embedded human-stomach-cancer, antibody was diluted at 1:200




Immunohistochemical analysis of paraffin-embedded Human skin. 1, Antibody was diluted at 1:200 (4°,overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .




Immunohistochemical analysis of paraffin-embedded Human skin. 1, Antibody was diluted at 1:200 (4°,overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .



Immunohistochemical analysis of paraffin-embedded Human skin. 1, Antibody was diluted at 1:200 (4°,overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .



Immunohistochemical analysis of paraffin-embedded Human gastric cancer. 1, Antibody was diluted at 1:100 (4°,overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .



Immunohistochemical analysis of paraffin-embedded Human gastric cancer. 1, Antibody was diluted at 1:100 (4°,overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .

## EnkiLife 优势产品推荐

| 产品类别                        | 产品货号     | 产品名称                                               |
|-----------------------------|----------|----------------------------------------------------|
| WB 解决方案                     | RA10020  | 2h 极速 WB 即用型全流程试剂盒                                 |
|                             | RA10021  | 4h 快速 WB 即用型全流程试剂盒                                 |
|                             | RA10042  | 5h 畅享版 WB 全流程试剂盒                                   |
|                             | RA10037  | 校准级彩色预染蛋白 Marker (8-180kDa)                        |
|                             | RA10038  | 校准级彩色预染蛋白 Marker (10-250kDa)                       |
|                             | RA10039  | 校准级高分子彩色预染蛋白 Marker (25-400kDa)                    |
| TSA 多重荧光染色试剂盒               | RA10008  | TSA 双标三色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10009  | TSA 三标四色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10010  | TSA 四标五色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10011  | TSA 五标六色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10012  | TSA 六标七色多重荧光染色试剂盒 (mIHC)                           |
| IHC 检测试剂盒                   | RA10006  | HRP Anti-Mouse/Rabbit IHC Detection System         |
|                             | RA10007  | Polymer-HRP Anti-Mouse/Rabbit IHC Detection System |
| 抗体标记试剂盒                     | RE80004p | 辣根过氧化物酶(HRP)抗体标记试剂盒                                |
|                             | RE80002q | Sulfo-NHS-生物素标记试剂盒                                 |
|                             | RE80007p | Cy3 荧光素标记试剂盒                                       |
|                             | RE80011p | Fluor488 荧光素标记试剂盒                                  |
|                             | RE80017p | Fluor750 荧光素标记试剂盒                                  |
|                             | RE80005p | 藻红蛋白(R-PE) 抗体快速标记试剂盒                               |
|                             | RE80040  | PE-Cy7 串联染料抗体快速标记试剂盒                               |
| 稳转细胞系构建服务<br>(免费赠送全膜 WB 验证) | TS-0001  | 过表达稳转细胞系构建                                         |
|                             | TS-0002  | 敲低稳转细胞系构建                                          |
|                             | TS-0003  | 敲除细胞系构建                                            |

限于篇幅，更多产品详见官网：<https://www.enkilife.cn>