

**产品名称: eIF3 $\alpha$  Rabbit Polyclonal Antibody**  
**产品货号: AP Rab10378**



## 产品概述 (Summary)

|                        |                                          |
|------------------------|------------------------------------------|
| 产品名称 (Production Name) | eIF3 $\alpha$ Rabbit Polyclonal Antibody |
| 描述 (Description)       | Rabbit polyclonal Antibody               |
| 宿主 (Host)              | Rabbit                                   |
| 应用 (Application)       | WB,ELISA                                 |
| 种属反应性 (Reactivity)     | Human,Mouse,Rat                          |

## 产品性能 (Performance)

|                     |                                                                                                   |
|---------------------|---------------------------------------------------------------------------------------------------|
| 偶联物 (Conjugation)   | Unconjugated                                                                                      |
| 修饰 (Modification)   | Unmodified                                                                                        |
| 同种型 (Isotype)       | IgG                                                                                               |
| 克隆 (Clonality)      | Polyclonal                                                                                        |
| 形式 (Form)           | Liquid                                                                                            |
| 存放说明 (Storage)      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.          |
| 储存溶液 (Buffer)       | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| 纯化方式 (Purification) | Affinity purification                                                                             |

## 免疫原信息 (Immunogen)

|                        |                                                                                                                                                         |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------|
| 基因名 (Gene Name)        | EIF3J                                                                                                                                                   |
| 别名 (Alternative Names) | EIF3J; EIF3S1; Eukaryotic translation initiation factor 3 subunit J; eIF3j; Eukaryotic translation initiation factor 3 subunit 1; eIF-3-alpha; eIF3 p35 |
| 基因 ID (Gene ID)        | 8669.0                                                                                                                                                  |
| 蛋白 ID (SwissProt ID)   | O75822.Synthesized peptide derived from eIF3 $\alpha$ . at AA range: 40-120                                                                             |

## 产品应用 (Application)

|                          |                                       |
|--------------------------|---------------------------------------|
| 稀释比 (Dilution Ratio)     | WB 1:500-1:2000,ELISA 1:20000-1:40000 |
| 蛋白分子量 (Molecular Weight) | 30kDa                                 |

## 研究背景 (Background)

**产品名称: eIF3 $\alpha$  Rabbit Polyclonal Antibody**  
**产品货号: AP Rab10378**




This gene encodes a core subunit of the eukaryotic initiation factor 3 complex, which participates in the initiation of translation by aiding in the recruitment of protein and mRNA components to the 40S ribosome. There are pseudogenes for this gene on chromosomes 1, 3, and 9. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Sep 2013],function:Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA<sub>i</sub> and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. This subunit binds directly within the mRNA entry channel of the 40S ribosome to the aminoacyl (A) site. It may regulate the interaction between the 43S PIC and mRNA.,mass spectrometry: PubMed:17322308,mass spectrometry:

PubMed:18599441,PTM:Phosphorylated. Phosphorylation is enhanced upon serum stimulation.,similarity:Belongs to the eIF-3 subunit J family.,subunit:Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is composed of 13 subunits: EIF3A, EIF3B, EIF3C, EIF3D, EIF3E, EIF3F, EIF3G, EIF3H, EIF3I, EIF3J, EIF3K, EIF3L and EIF3M. The eIF-3 complex appears to include 3 stable modules: module A is composed of EIF3A, EIF3B, EIF3G and EIF3I; module B is composed of EIF3F, EIF3H, and EIF3M; and module C is composed of EIF3C, EIF3D, EIF3E, EIF3K and EIF3L. EIF3C of module C binds EIF3B of module A and EIF3H of module B, thereby linking the three modules. EIF3J is a labile subunit that binds to the eIF-3 complex via EIF3B. The eIF-3 complex interacts with RPS6KB1 under conditions of nutrient depletion. Mitogenic stimulation leads to binding and activation of a complex composed of FRAP1 and RAPTOR, leading to phosphorylation and release of RPS6KB1 and binding of EIF4B to eIF-3.,

## 研究领域 (Research Area)

## 图片 (Image Data)



Western Blot analysis of various cells using Antibody diluted at 1:1000. Secondary antibody was diluted at 1:20000

**产品名称: eIF3 $\alpha$  Rabbit Polyclonal Antibody**  
**产品货号: APRab10378**



---

**注意事项 (Note)**

For research use only .