产品名称：Endophilin I Rabbit Polyclonal Antibody
产品货号：APRab10467

产品概述（Summary）

产品名称（Production Name）	Endophilin I Rabbit Polyclonal Antibody
描述（Description）	Rabbit polyclonal Antibody
宿主（Host）	Rabbit
应用（Application）	WB,ELISA
种属反应性（Reactivity）	Human,Mouse,Rat

产品性能（Performance）

偶联物（Conjugation）	Unconjugated
修饰（Modification）	Unmodified
同种型（Isotype）	IgG
克隆（Clonality）	Polyclonal
形式（Form）	Liquid
存放说明（Storage）	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液（Buffer）	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式（Purification）	Affinity purification

免疫原信息（Immunogen）

基因名（Gene Name）	SH3GL2
别名（Alternative Names）	SH3GL2; CNSA2; SH3D2A; Endophilin-A1; EEN-B1; Endophilin-1; SH3 domain protein 2A; SH3 domain-containing GRB2-like protein 2
基因 ID（Gene ID）	6456.0
蛋白 ID（SwissProt ID）	Q99962.Synthesized peptide derived from Endophilin I . at AA range: 30-110

产品应用（Application）

稀释比（Dilution Ratio）	WB 1:500-1:2000,ELISA 1:5000-1:20000
蛋白分子量（Molecular Weight）	39kDa

研究背景（Background）

domain:An N-terminal amphipathic helix, the BAR domain and a second amphipathic helix inserted into helix 1 of the BAR domain (N-BAR domain) induce membrane curvature and bind curved membranes. The BAR domain dimer forms a rigid crescent shaped bundle of helices with the pair of second amphipathic helices protruding towards the membrane-binding surface.,function:Implicated in synaptic vesicle endocytosis. May recruit other proteins to membranes with high curvature.,miscellaneous:HeLa cells expressing the N-BAR domain of SH3GL2 show tubulation of the plasma membrane. The N-BAR domain binds liposomes and induces formation of tubules from liposomes. The N-terminal amphipathic helix is required for liposome binding. The second amphipathic helix enhances liposome tubulation.,similarity:Belongs to the endophilin family.,similarity:Contains 1 BAR domain.,similarity:Contains 1 SH3 domain.,subcellular location:Concentrated in presynaptic nerve terminals in neurons.,subunit:Monomer; in cytoplasm. Homodimer; when associated with membranes (By similarity). Interacts with SYNJ1 and DNM1. Interacts with MAP4K3; the interaction appears to regulate MAP4K3-mediated JNK activation. Interacts with PDCD6IP.,tissue specificity:Brain, mostly in frontal cortex. Expressed at high level in fetal cerebellum.,domain:An N-terminal amphipathic helix, the BAR domain and a second amphipathic helix inserted into helix 1 of the BAR domain (N-BAR domain) induce membrane curvature and bind curved membranes. The BAR domain dimer forms a rigid crescent shaped bundle of helices with the pair of second amphipathic helices protruding towards the membrane-binding surface.,function:Implicated in synaptic vesicle endocytosis. May recruit other proteins to membranes with high curvature.,miscellaneous:HeLa cells expressing the N-BAR domain of SH3GL2 show tubulation of the plasma membrane. The N-BAR domain binds liposomes and induces formation of tubules from liposomes. The N-terminal amphipathic helix is required for liposome binding. The second amphipathic helix enhances liposome tubulation.,similarity:Belongs to the endophilin family.,similarity:Contains 1 BAR domain.,similarity:Contains 1 SH3 domain.,subcellular location:Concentrated in presynaptic nerve terminals in neurons.,subunit:Monomer; in cytoplasm. Homodimer; when associated with membranes (By similarity). Interacts with SYNJ1 and DNM1. Interacts with MAP4K3; the interaction appears to regulate MAP4K3-mediated JNK activation. Interacts with PDCD6IP.,tissue specificity:Brain, mostly in frontal cortex. Expressed at high level in fetal cerebellum.,

研究领域（Research Area）

Endocytosis;

图片（Image Data）

产品名称：Endophilin I Rabbit Polyclonal Antibody
产品货号：APRab10467



Western blot analysis of SH3GL2 Antibody. The lane on the right is blocked with the SH3GL2 peptide.

注意事项（Note）

For research use only .