

产品名称: Enterokinase HC Rabbit Polyclonal Antibody
产品货号: APRab10482



产品概述 (Summary)

产品名称 (Production Name)	Enterokinase HC Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Rat, Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	TMPRSS15
别名 (Alternative Names)	TMPRSS15; ENTK; PRSS7; Enteropeptidase; Enterokinase; Serine protease 7; Transmembrane protease serine 15
基因 ID (Gene ID)	5651.0
蛋白 ID (SwissProt ID)	P98073. The antiserum was produced against synthesized peptide derived from human ENTK. AA range: 81-130

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:100-1:300, ICC/IF 1:200-1:1000, ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	

产品名称: Enterokinase HC Rabbit Polyclonal Antibody
产品货号: APRab10482




研究背景 (Background)

This gene encodes an enzyme that converts the pancreatic proenzyme trypsinogen to trypsin, which activates other proenzymes including chymotrypsinogen and procarboxypeptidases. The precursor protein is cleaved into two chains that form a heterodimer linked by a disulfide bond. This protein is a member of the trypsin family of peptidases. Mutations in this gene cause enterokinase deficiency, a malabsorption disorder characterized by diarrhea and failure to thrive. [provided by RefSeq, Jul 2008],catalytic activity:Activation of trypsinogen by selective cleavage of 6-Lys-Ile-7 bond.,disease:Defects in PRSS7 are a cause of enterokinase deficiency [MIM:226200]; a life-threatening intestinal malabsorption disorder characterized by diarrhea and failure to thrive.,function:Responsible for initiating activation of pancreatic proteolytic proenzymes (trypsin, chymotrypsin and carboxypeptidase A). It catalyzes the conversion of trypsinogen to trypsin which in turn activates other proenzymes including chymotrypsinogen, procarboxypeptidases, and proelastases.,PTM:The chains are derived from a single precursor that is cleaved by a trypsin-like protease.,similarity:Belongs to the peptidase S1 family.,similarity:Contains 1 MAM domain.,similarity:Contains 1 peptidase S1 domain.,similarity:Contains 1 SEA domain.,similarity:Contains 1 SRCR domain.,similarity:Contains 2 CUB domains.,similarity:Contains 2 LDL-receptor class A domains.,subunit:Heterodimer of a catalytic (light) chain and a multidomain (heavy) chain linked by a disulfide bond.,tissue specificity:Intestinal brush border.,

研究领域 (Research Area)

图片 (Image Data)



Immunofluorescence analysis of HepG2 cells, using ENTK Antibody. The picture on the right is blocked with the synthesized peptide.

注意事项 (Note)

For research use only .