

产品名称: EWS Rabbit Polyclonal Antibody

产品货号: APRab10653

产品概述 (Summary)

描述(Description)	Rabbit polyclonal Antibody
宿主(Host)	Rabbit
应用(Application)	WB,ELISA
种属反应(Reactivity)	Human,Mouse,Rat
偶联物(Conjugation)	Unconjugated
修饰(Modification)	Unmodified
同种型(Isotype)	IgG
克隆(Clonality)	Polyclonal
剂型(Form)	Liquid
保存条件(Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液(Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化 (Purification)	Affinity purification

产品应用(Application)

稀释比(Dilution Ratio)	WB 1:500-1:2000,ELISA 1:5000-1:20000
分子量(Molecular Weight)	68kDa

抗原信息(Antigen Information)

基因名(Gene Name)	EWSR1
别名(Alternative Names)	EWSR1; EWS; RNA-binding protein EWS; EWS oncogene; Ewing sarcoma breakpoint region 1 protein
基因 ID(Gene ID)	2130.0
SwissProt ID	Q01844
免疫原(Immunogen)	The antiserum was produced against synthesized peptide derived from human EWSR1. AA range:403-452

研究背景 (Background)

This gene encodes a multifunctional protein that is involved in various cellular processes, including gene expression, cell

signaling, and RNA processing and transport. The protein includes an N-terminal transcriptional activation domain and a C-terminal RNA-binding domain. Chromosomal translocations between this gene and various genes encoding transcription factors result in the production of chimeric proteins that are involved in tumorigenesis. These chimeric proteins usually consist of the N-terminal transcriptional activation domain of this protein fused to the C-terminal DNA-binding domain of the transcription factor protein. Mutations in this gene, specifically a t(11;22)(q24;q12) translocation, are known to cause Ewing sarcoma as well as neuroectodermal and various other tumors. Alternative splicing of this gene results in multiple transcript variants. Related pseudogenes have been identified.

disease:A chromosomal aberration involving EWSR1 is associated with desmoplastic small round cell tumor (DSRCT). Translocation t(11;22)(p13;q12) with WT1.

disease:A chromosomal aberration involving EWSR1 is associated with malignant melanoma of soft parts (MMSP). Translocation t(12;22)(q13;q12) with ATF-1. Malignant melanoma of soft parts, also known as soft tissue clear cell sarcoma, is a rare tumor developing in tendons and aponeuroses.

disease:A chromosomal aberration involving EWSR1 is associated with small round cell sarcoma. Translocation t(11;22)(p36.1;q12) with PATZ1.

disease:Chromosomal aberrations involving EWSR1 are a cause of Ewing sarcoma [MIM:133450]. Translocation t(11;22)(q24;q12) with FLI1; translocation t(7;22)(p22;q12) with ETV1; translocation t(21;22)(q22;q12) with ERG; translocation t(9;22)(q22-31;q11-12) with NR4A3. Translocation t(2;21;22)(q23;q22;q12) that forms a EWSR1-FEV fusion protein with potential oncogenic activity.

disease:Chromosomal aberrations involving EWSR1 are associated with angiomatoid fibrous histiocytoma (AFH) [MIM:612160]. Translocation t(12;22)(q13;q12) with ATF1 generates a chimeric EWSR1/ATF1 protein. Translocation t(2;22)(q33;q12) with CREB1 generates a EWSR1/CREB1 fusion gene that is most common genetic abnormality in this tumor type.

domain:EWS activation domain (EAD) functions as a potent activation domain in EFPS. EWSR1 binds POLR2C but not POLR2E or POLR2G, whereas the isolated EAD binds POLR2E and POLR2G but not POLR2C. Cis-linked RNA-binding domain (RBD) can strongly and specifically repress trans-activation by the EAD.

function:Might normally function as a repressor. EWS-fusion-proteins (EFPS) may play a role in the tumorigenic process. They may disturb gene expression by mimicking, or interfering with the normal function of CTD-POLII within the transcription initiation complex. They may also contribute to an aberrant activation of the fusion protein target genes.

miscellaneous:Binds calmodulin in the presence, but not in the absence, of calcium ion.

miscellaneous:EFPS arise due to chromosomal translocations in which EWSR1 is fused to a variety of cellular transcription factors. EFPS are very potent transcriptional activators dependent on the EAD and a C-terminal DNA-binding domain contributed by the fusion partner. The spectrum of malignancies associated with EFPS are thought to arise via EFP-induced transcriptional deregulation, with the tumor phenotype specified by the EWSR1 fusion partner and cell type. Transcriptional repression of the transforming growth factor beta type II receptor (TGF beta RII) is an important target of the EWS-FLI1, EWS-ERG, or EWS-ETV1 oncogene.

PTM:Highly methylated on arginine residues. Methylation is mediated by PRMT1 and, at lower level by PRMT8.

PTM:Phosphorylated; calmodulin-binding inhibits phosphorylation of Ser-266.

similarity:Belongs to the ETS family.

similarity:Belongs to the RRM TET family.

similarity:Contains 1 ETS DNA-binding domain.

similarity:Contains 1 IQ domain.

similarity:Contains 1 RanBP2-type zinc finger.

similarity:Contains 1 RRM (RNA recognition motif) domain.

subcellular location:Relocates from cytoplasm to ribosomes upon PTK2B/FAK2 activation.

subunit:Binds POLR2C, SF1, calmodulin and RNA. Interacts with PTK2B/FAK2 and TDRD3.

tissue specificity:Ubiquitous.


研究领域 (Research Area)

Tags & Cell Markers


注意事项 (Note)

For research use only.


图片 (Image Data)



Western blot analysis of the lysates from HepG2 cells using EWSR1 antibody.



Western Blot analysis of various cells using EWS Polyclonal Antibody



Western Blot analysis of COLO205 cells using EWS Polyclonal Antibody

EnkiLife 优势产品推荐

产品类别	产品货号	产品名称
WB 解决方案	RA10020	2h 极速 WB 即用型全流程试剂盒
	RA10021	4h 快速 WB 即用型全流程试剂盒
	RA10042	5h 畅享版 WB 全流程试剂盒
	RA10037	校准级彩色预染蛋白 Marker (8-180kDa)
	RA10038	校准级彩色预染蛋白 Marker (10-250kDa)
	RA10039	校准级高分子彩色预染蛋白 Marker (25-400kDa)
TSA 多重荧光染色试剂盒	RA10008	TSA 双标三色多重荧光染色试剂盒 (mIHC)
	RA10009	TSA 三标四色多重荧光染色试剂盒 (mIHC)
	RA10010	TSA 四标五色多重荧光染色试剂盒 (mIHC)
	RA10011	TSA 五标六色多重荧光染色试剂盒 (mIHC)
	RA10012	TSA 六标七色多重荧光染色试剂盒 (mIHC)
IHC 检测试剂盒	RA10006	HRP Anti-Mouse/Rabbit IHC Detection System
	RA10007	Polymer-HRP Anti-Mouse/Rabbit IHC Detection System
抗体标记试剂盒	RE80004p	辣根过氧化物酶(HRP)抗体标记试剂盒
	RE80002q	Sulfo-NHS-生物素标记试剂盒
	RE80007p	Cy3 荧光素标记试剂盒
	RE80011p	Fluor488 荧光素标记试剂盒
	RE80017p	Fluor750 荧光素标记试剂盒
	RE80005p	藻红蛋白(R-PE) 抗体快速标记试剂盒
	RE80040	PE-Cy7 串联染料抗体快速标记试剂盒
稳转细胞系构建服务 (免费赠送全膜 WB 验证)	TS-0001	过表达稳转细胞系构建
	TS-0002	敲低稳转细胞系构建
	TS-0003	敲除细胞系构建

限于篇幅，更多产品详见官网：<https://www.enkilife.cn>