

产品名称: FAP-1 Rabbit Polyclonal Antibody
产品货号: AP Rab10831



产品概述 (Summary)

产品名称 (Production Name)	FAP-1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	FAP
别名 (Alternative Names)	Seprase (EC 3.4.21.-) (170 kDa melanoma membrane-bound gelatinase) (Fibroblast activation protein alpha) (Integral membrane serine protease)
基因 ID (Gene ID)	2191.0
蛋白 ID (SwissProt ID)	Q12884.Synthesized peptide derived from human FAP-1 Polyclonal

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	90kDa

研究背景 (Background)

产品名称: FAP-1 Rabbit Polyclonal Antibody
产品货号: AP Rab10831




The protein encoded by this gene is a homodimeric integral membrane gelatinase belonging to the serine protease family. It is selectively expressed in reactive stromal fibroblasts of epithelial cancers, granulation tissue of healing wounds, and malignant cells of bone and soft tissue sarcomas. This protein is thought to be involved in the control of fibroblast growth or epithelial-mesenchymal interactions during development, tissue repair, and epithelial carcinogenesis. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Apr 2014], catalytic activity: Degrades gelatin and heat-denatured type I and type IV collagen, but not native type I or type IV collagen. Does not cleave laminin, fibronectin, fibrin or casein., function: May have a role in tissue remodeling during development and wound healing, and may contribute to invasiveness in malignant cancers., induction: In fibroblasts at times and sites of tissue remodeling during development, tissue repair, and carcinogenesis., PTM: N-glycosylated., PTM: The N-terminus may be blocked., similarity: Belongs to the peptidase S9B family., subcellular location: Found in cell surface lamellipodia, invadopodia and on shed vesicles., subunit: Homodimer, or heterodimer with DPP4. The monomer is inactive., tissue specificity: Fibroblast specific.,

研究领域 (Research Area)

Cell Biology

图片 (Image Data)



Western blot analysis of mouse-lung lysate, antibody was diluted at 1000. Secondary antibody was diluted at 1:20000

注意事项 (Note)

For research use only .