

产品名称: Fish Rabbit Polyclonal Antibody

产品货号: APRab10993

产品概述 (Summary)

描述(Description)	Rabbit polyclonal Antibody
宿主(Host)	Rabbit
应用(Application)	WB,IHC,ICC/IF,ELISA
种属反应(Reactivity)	Human,Mouse
偶联物(Conjugation)	Unconjugated
修饰(Modification)	Unmodified
同种型(Isotype)	IgG
克隆(Clonality)	Polyclonal
剂型(Form)	Liquid
保存条件(Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液(Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化 (Purification)	Affinity purification

产品应用(Application)

稀释比(Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:500,ICC/IF 1:100-1:500,ELISA 1:5000-1:20000

分子量(Molecular Weight)

抗原信息(Antigen Information)

基因名(Gene Name)	SH3PXD2A SH3PXD2A; FISH; KIAA0418; SH3MD1; TKS5; SH3 and PX domain-containing protein 2A;
别名(Alternative Names)	Adapter protein TKS5; Five SH3 domain-containing protein; SH3 multiple domains protein 1; Tyrosine kinase substrate with five SH3 domains
基因 ID(Gene ID)	9644.0
SwissProt ID	Q5TCZ1
免疫原(Immunogen)	Synthesized peptide derived from the N-terminal region of human Fish.

研究背景 (Background)

domain:The fifth SH3 domain mediates binding with ADAM12, ADAM15 and ADAM19.,domain:The PX domain is required for


podosome localization, and for binding phosphatidylinositol 3-phosphate (PtdIns(3)P) and phosphatidylinositol 3,4-bisphosphate (PtdIns(3,4)P₂).,function:Required for podosome formation, degradation of the extracellular matrix, and for the invasiveness of some cancer cells. Binds phosphatidylinositol 3-phosphate (PtdIns(3)P) and phosphatidylinositol 3,4-bisphosphate (PtdIns(3,4)P₂). In association with ADAM12, mediates the neurotoxic effect of beta-amyloid peptide.,PTM:Tyrosine phosphorylated by SRC. Phosphorylation plays a regulatory role in the protein localization. The intramolecular interaction of the PX domain with the third SH3 domain maintains the protein in the cytoplasm and phosphorylation disrupts this interaction, resulting in the redistribution of the protein from cytoplasm to the perimembrane region. Phosphorylated on serine upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the SH3PXD2 family.,similarity:Contains 1 PX (phox homology) domain.,similarity:Contains 5 SH3 domains.,subcellular location:Cytoplasmic in normal cells and localizes to podosomes in SRC-transformed cells.,subunit:Interacts with ADAM12, ADAM15 and ADAM19.,tissue specificity:Found in several cancer cell lines, particularly invasive breast carcinomas and melanomas.,domain:The fifth SH3 domain mediates binding with ADAM12, ADAM15 and ADAM19.,domain:The PX domain is required for podosome localization, and for binding phosphatidylinositol 3-phosphate (PtdIns(3)P) and phosphatidylinositol 3,4-bisphosphate (PtdIns(3,4)P₂).,function:Required for podosome formation, degradation of the extracellular matrix, and for the invasiveness of some cancer cells. Binds phosphatidylinositol 3-phosphate (PtdIns(3)P) and phosphatidylinositol 3,4-bisphosphate (PtdIns(3,4)P₂). In association with ADAM12, mediates the neurotoxic effect of beta-amyloid peptide.,PTM:Tyrosine phosphorylated by SRC. Phosphorylation plays a regulatory role in the protein localization. The intramolecular interaction of the PX domain with the third SH3 domain maintains the protein in the cytoplasm and phosphorylation disrupts this interaction, resulting in the redistribution of the protein from cytoplasm to the perimembrane region. Phosphorylated on serine upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the SH3PXD2 family.,similarity:Contains 1 PX (phox homology) domain.,similarity:Contains 5 SH3 domains.,subcellular location:Cytoplasmic in normal cells and localizes to podosomes in SRC-transformed cells.,subunit:Interacts with ADAM12, ADAM15 and ADAM19.,tissue specificity:Found in several cancer cell lines, particularly invasive breast carcinomas and melanomas.,

研究领域 (Research Area)

注意事项 (Note)

For research use only.

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human uterus. 1, Antibody was diluted at 1:200 (4° overnight) . 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min) .

EnkiLife 优势产品推荐

产品类别	产品货号	产品名称
WB 解决方案	RA10020	2h 极速 WB 即用型全流程试剂盒
	RA10021	4h 快速 WB 即用型全流程试剂盒
	RA10042	5h 畅享版 WB 全流程试剂盒
	RA10037	校准级彩色预染蛋白 Marker (8-180kDa)
	RA10038	校准级彩色预染蛋白 Marker (10-250kDa)
	RA10039	校准级高分子彩色预染蛋白 Marker (25-400kDa)
TSA 多重荧光染色试剂盒	RA10008	TSA 双标三色多重荧光染色试剂盒 (mIHC)
	RA10009	TSA 三标四色多重荧光染色试剂盒 (mIHC)
	RA10010	TSA 四标五色多重荧光染色试剂盒 (mIHC)
	RA10011	TSA 五标六色多重荧光染色试剂盒 (mIHC)
	RA10012	TSA 六标七色多重荧光染色试剂盒 (mIHC)
IHC 检测试剂盒	RA10006	HRP Anti-Mouse/Rabbit IHC Detection System
	RA10007	Polymer-HRP Anti-Mouse/Rabbit IHC Detection System
抗体标记试剂盒	RE80004p	辣根过氧化物酶(HRP)抗体标记试剂盒
	RE80002q	Sulfo-NHS-生物素标记试剂盒
	RE80007p	Cy3 荧光素标记试剂盒
	RE80011p	Fluor488 荧光素标记试剂盒
	RE80017p	Fluor750 荧光素标记试剂盒
	RE80005p	藻红蛋白(R-PE) 抗体快速标记试剂盒
	RE80040	PE-Cy7 串联染料抗体快速标记试剂盒
稳转细胞系构建服务 (免费赠送全膜 WB 验证)	TS-0001	过表达稳转细胞系构建
	TS-0002	敲低稳转细胞系构建
	TS-0003	敲除细胞系构建

限于篇幅，更多产品详见官网：<https://www.enkilife.cn>