

产品名称: FoxO1/3 Rabbit Polyclonal Antibody
产品货号: APRab11099



产品概述 (Summary)

产品名称 (Production Name)	FoxO1/3 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	FOXO1/FOXO3 FOXO1; FKHR; FOXO1A; Forkhead box protein O1; Forkhead box protein O1A;
别名 (Alternative Names)	Forkhead in rhabdomyosarcoma; FOXO3; FKHL1; FOXO3A; Forkhead box protein O3; AF6q21 protein; Forkhead in rhabdomyosarcoma-like 1
基因 ID (Gene ID)	2308.0
蛋白 ID (SwissProt ID)	Q12778/O43524. The antiserum was produced against synthesized peptide derived from human FOXO1A/3A. AA range: 291-340

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	

产品名称: FoxO1/3 Rabbit Polyclonal Antibody
产品货号: AP Rab11099




研究背景 (Background)

This gene belongs to the forkhead family of transcription factors which are characterized by a distinct forkhead domain. The specific function of this gene has not yet been determined; however, it may play a role in myogenic growth and differentiation. Translocation of this gene with PAX3 has been associated with alveolar rhabdomyosarcoma. [provided by RefSeq, Jul 2008], disease: Chromosomal aberrations involving FOXO1 are a cause of rhabdomyosarcoma 2 (RMS2) [MIM:268220]; also known as alveolar rhabdomyosarcoma. Translocation (2;13)(q35;q14) with PAX3; translocation t(1;13)(p36;q14) with PAX7. The resulting protein is a transcriptional activator., function: Transcription factor., PTM: Phosphorylated by AKT1; insulin-induced (By similarity). IGF1 rapidly induces phosphorylation of Ser-256, Thr-24, and Ser-319. Phosphorylation of Ser-256 decreases DNA-binding activity and promotes the phosphorylation of Thr-24, and Ser-319, permitting phosphorylation of Ser-322 and Ser-325, probably by CK1, leading to nuclear exclusion and loss of function. Phosphorylation of Ser-329 is independent of IGF1 and leads to reduced function. Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Contains 1 fork-head DNA-binding domain., subcellular location: Shuttles between cytoplasm and nucleus., subunit: Interacts with LRPPRC., tissue specificity: Ubiquitous.,

研究领域 (Research Area)

Insulin Receptor; B Cell Receptor; Protein_Acetylation

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human thyroid gland tissue, using FOXO1A/3A Antibody. The picture on the right is blocked with the synthesized peptide.

注意事项 (Note)

For research use only .