

**产品名称:** Frizzled-8 Rabbit Polyclonal Antibody

**产品货号:** APRab11149

### 产品概述 (Summary)

|                   |                                                                                                   |
|-------------------|---------------------------------------------------------------------------------------------------|
| 描述(Description)   | Rabbit polyclonal Antibody                                                                        |
| 宿主(Host)          | Rabbit                                                                                            |
| 应用(Application)   | WB,IHC,ICC/IF,ELISA                                                                               |
| 种属反应(Reactivity)  | Human,Mouse,Rat                                                                                   |
| 偶联物(Conjugation)  | Unconjugated                                                                                      |
| 修饰(Modification)  | Unmodified                                                                                        |
| 同种型(Isotype)      | IgG                                                                                               |
| 克隆(Clonality)     | Polyclonal                                                                                        |
| 剂型(Form)          | Liquid                                                                                            |
| 保存条件(Storage)     | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.          |
| 储存溶液(Buffer)      | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| 纯化 (Purification) | Affinity purification                                                                             |

### 产品应用(Application)

|                       |                                                                        |
|-----------------------|------------------------------------------------------------------------|
| 稀释比(Dilution Ratio)   | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:10000 |
| 分子量(Molecular Weight) | 70kDa                                                                  |

### 抗原信息(Antigen Information)

|                       |                                                                                                  |
|-----------------------|--------------------------------------------------------------------------------------------------|
| 基因名(Gene Name)        | FZD8                                                                                             |
| 别名(Alternative Names) | FZD8; Frizzled-8; Fz-8; hFz8                                                                     |
| 基因 ID(Gene ID)        | 8325.0                                                                                           |
| SwissProt ID          | Q9H461                                                                                           |
| 免疫原(Immunogen)        | The antiserum was produced against synthesized peptide derived from human FZD8. AA range:486-535 |

### 研究背景 (Background)

frizzled class receptor 8(FZD8) Homo sapiens This intronless gene is a member of the frizzled gene family. Members of this family encode seven-transmembrane domain proteins that are receptors for the Wntless type MMTV integration site family of

signaling proteins. Most frizzled receptors are coupled to the beta-catenin canonical signaling pathway. This gene is highly expressed in two human cancer cell lines, indicating that it may play a role in several types of cancer. The crystal structure of the extracellular cysteine-rich domain of a similar mouse protein has been determined. [provided by RefSeq, Jul 2008],domain:Lys-Thr-X-X-X-Trp motif is involved in the activation of the Wnt/beta-catenin signaling pathway.,domain:The FZ domain is involved in binding with Wnt ligands.,domain:The PDZ-binding motif mediates interaction with GOPC.,function:Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues.,similarity:Belongs to the G-protein coupled receptor Fz/Smo family.,similarity:Contains 1 FZ (frizzled) domain.,subunit:Interacts with GOPC. Interacts with RSPO1 and RSPO3.,tissue specificity:Most abundant in fetal kidney, followed by brain and lung. In adult tissues, expressed in kidney, heart, pancreas and skeletal muscle.,


### 研究领域 (Research Area)

WNT;WNT-T CELLMelanogenesis;Pathways in cancer;Colorectal cancer;Basal cell carcinoma;

### 注意事项 (Note)

For research use only.

### 图片 (Image Data)



Western blot analysis of lysates from Jurkat cells, using FZD8 Antibody. The lane on the right is blocked with the synthesized peptide.

## EnkiLife 优势产品推荐

| 产品类别                        | 产品货号     | 产品名称                                               |
|-----------------------------|----------|----------------------------------------------------|
| WB 解决方案                     | RA10020  | 2h 极速 WB 即用型全流程试剂盒                                 |
|                             | RA10021  | 4h 快速 WB 即用型全流程试剂盒                                 |
|                             | RA10042  | 5h 畅享版 WB 全流程试剂盒                                   |
|                             | RA10037  | 校准级彩色预染蛋白 Marker (8-180kDa)                        |
|                             | RA10038  | 校准级彩色预染蛋白 Marker (10-250kDa)                       |
|                             | RA10039  | 校准级高分子彩色预染蛋白 Marker (25-400kDa)                    |
| TSA 多重荧光染色试剂盒               | RA10008  | TSA 双标三色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10009  | TSA 三标四色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10010  | TSA 四标五色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10011  | TSA 五标六色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10012  | TSA 六标七色多重荧光染色试剂盒 (mIHC)                           |
| IHC 检测试剂盒                   | RA10006  | HRP Anti-Mouse/Rabbit IHC Detection System         |
|                             | RA10007  | Polymer-HRP Anti-Mouse/Rabbit IHC Detection System |
| 抗体标记试剂盒                     | RE80004p | 辣根过氧化物酶(HRP)抗体标记试剂盒                                |
|                             | RE80002q | Sulfo-NHS-生物素标记试剂盒                                 |
|                             | RE80007p | Cy3 荧光素标记试剂盒                                       |
|                             | RE80011p | Fluor488 荧光素标记试剂盒                                  |
|                             | RE80017p | Fluor750 荧光素标记试剂盒                                  |
|                             | RE80005p | 藻红蛋白(R-PE) 抗体快速标记试剂盒                               |
|                             | RE80040  | PE-Cy7 串联染料抗体快速标记试剂盒                               |
| 稳转细胞系构建服务<br>(免费赠送全膜 WB 验证) | TS-0001  | 过表达稳转细胞系构建                                         |
|                             | TS-0002  | 敲低稳转细胞系构建                                          |
|                             | TS-0003  | 敲除细胞系构建                                            |

限于篇幅，更多产品详见官网：<https://www.enkilife.cn>