

产品名称: Frizzled-9 Rabbit Polyclonal Antibody
产品货号: APRab11150



产品概述 (Summary)

产品名称 (Production Name)	Frizzled-9 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Monkey

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	FZD9
别名 (Alternative Names)	FZD9; FZD3; Frizzled-9; Fz-9; hFz9; FzE6; CD antigen CD349
基因 ID (Gene ID)	8326.0
蛋白 ID (SwissProt ID)	O00144.The antiserum was produced against synthesized peptide derived from human FZD9. AA range:542-591

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	64kDa

研究背景 (Background)

产品名称: Frizzled-9 Rabbit Polyclonal Antibody
产品货号: APRab11150




frizzled class receptor 9(FZD9) Homo sapiens Members of the 'frizzled' gene family encode 7-transmembrane domain proteins that are receptors for Wnt signaling proteins. The FZD9 gene is located within the Williams syndrome common deletion region of chromosome 7, and heterozygous deletion of the FZD9 gene may contribute to the Williams syndrome phenotype. FZD9 is expressed predominantly in brain, testis, eye, skeletal muscle, and kidney. [provided by RefSeq, Jul 2008],caution:Has been first described as FZD3 in litterature.,domain:Lys-Thr-X-X-X-Trp motif is involved in the activation of the Wnt/beta-catenin signaling pathway.,domain:The FZ domain is involved in binding with Wnt ligands.,function:Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues.,similarity:Belongs to the G-protein coupled receptor Fz/Smo family.,similarity:Contains 1 FZ (frizzled) domain.,tissue specificity:Expressed predominantly in adult and fetal brain, testis, eye, skeletal muscle and kidney. Moderately expressed in pancreas, thyroid, adrenal cortex, small intestine and stomach. Detected in fetal liver and kidney.,

研究领域 (Research Area)


WNT;WNT-T CELLMelanogenesis;Pathways in cancer;Colorectal cancer;Basal cell carcinoma;

图片 (Image Data)




Immunofluorescence analysis of A549 cells, using FZD9 Antibody. The picture on the right is blocked with the synthesized peptide.


产品名称: Frizzled-9 Rabbit Polyclonal Antibody
产品货号: APRab11150



Western blot analysis of lysates from HUVEC, COLO, and COS cells, using FZD9 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from Jurkat cells using FZD9 antibody.



Western Blot analysis of various cells using Frizzled-9 Polyclonal Antibody

注意事项 (Note)

For research use only .