

产品名称: Gastrin Rabbit Polyclonal Antibody
产品货号: APRab11307



产品概述 (Summary)

产品名称 (Production Name)	Gastrin Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	GAST
别名 (Alternative Names)	GAST; GAS; Gastrin
基因 ID (Gene ID)	2520.0
蛋白 ID (SwissProt ID)	P01350.The antiserum was produced against synthesized peptide derived from human Gastrin. AA range:52-101

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	12kDa

研究背景 (Background)


产品名称: Gastrin Rabbit Polyclonal Antibody
产品货号: AP Rab11307




Gastrin is a hormone whose main function is to stimulate secretion of hydrochloric acid by the gastric mucosa, which results in gastrin formation inhibition. This hormone also acts as a mitogenic factor for gastrointestinal epithelial cells. Gastrin has two biologically active peptide forms, G34 and G17. [provided by RefSeq, Jul 2008],function:Gastrin stimulates the stomach mucosa to produce and secrete hydrochloric acid and the pancreas to secrete its digestive enzymes. It also stimulates smooth muscle contraction and increases blood circulation and water secretion in the stomach and intestine.,online information:Gastrin entry,PTM:Sulfation enhances proteolytic processing, and blocks peptide degradation. Levels of sulfation differ between proteolytically-cleaved gastrins. Thus, gastrin-6 is almost 73% sulfated, whereas the larger gastrins are less than 50% sulfated. Sulfation levels are also tissue-specific.,PTM:Two different processing pathways probably exist in antral G-cells. In the dominant pathway progastrin is cleaved at three sites resulting in two major bioactive gastrins, gastrin-34 and gastrin-17. In the putative alternative pathway, progastrin may be processed only at the most C-terminal dibasic site resulting in the synthesis of gastrin-71.,similarity:Belongs to the gastrin/cholecystokinin family.,

研究领域 (Research Area)

图片 (Image Data)



Immunofluorescence analysis of HeLa cells, using Gastrin Antibody. The picture on the right is blocked with the synthesized peptide.




Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using Gastrin Antibody. The picture on

产品名称: Gastrin Rabbit Polyclonal Antibody
产品货号: APRab11307



the right is blocked with the synthesized peptide.



Western blot analysis of lysates from NIH/3T3 cells, using Gastrin Antibody. The lane on the right is blocked with the synthesized peptide.

注意事项 (Note)

For research use only .