

产品名称: GFR α -1 Rabbit Polyclonal Antibody
产品货号: APRab11418



产品概述 (Summary)

产品名称 (Production Name)	GFR α -1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	GFRA1 GFRA1; GDNFRA; RETL1; TRNR1; GDNF family receptor alpha-1; GDNF receptor alpha-1; GDNFR-alpha-1; GFR-alpha-1; RET ligand 1; TGF-beta-related neurotrophic factor receptor 1
别名 (Alternative Names)	
基因 ID (Gene ID)	2674.0
蛋白 ID (SwissProt ID)	P56159.The antiserum was produced against synthesized peptide derived from human GFR alpha-1. AA range:51-100

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	

产品名称: GFR α -1 Rabbit Polyclonal Antibody
产品货号: APRab11418




研究背景 (Background)

This gene encodes a member of the glial cell line-derived neurotrophic factor receptor (GDNFR) family of proteins. The encoded preproprotein is proteolytically processed to generate the mature receptor. Glial cell line-derived neurotrophic factor (GDNF) and neurturin (NTN) are two structurally related, potent neurotrophic factors that play key roles in the control of neuron survival and differentiation. This receptor is a glycosylphosphatidylinositol (GPI)-linked cell surface receptor for both GDNF and NTN, and mediates activation of the RET tyrosine kinase receptor. This gene is a candidate gene for Hirschsprung disease. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed. [provided by RefSeq, Jan 2016],function:Receptor for GDNF. Mediates the GDNF-induced autophosphorylation and activation of the RET receptor.,similarity:Belongs to the GDNFR family.,subunit:2 molecules of GDNFR-alpha are thought to form a complex with the disulfide-linked GDNF dimer and with 2 molecules of RET.,

研究领域 (Research Area)

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using GFR alpha-1 Antibody. The picture on the right is blocked with the synthesized peptide.

注意事项 (Note)

For research use only .