

产品名称: GLCNE Rabbit Polyclonal Antibody

产品货号: APRab11459

产品概述 (Summary)

描述(Description)	Rabbit polyclonal Antibody
宿主(Host)	Rabbit
应用(Application)	WB,IHC,ICC/IF,ELISA
种属反应(Reactivity)	Human,Mouse,Rat
偶联物(Conjugation)	Unconjugated
修饰(Modification)	Unmodified
同种型(Isotype)	IgG
克隆(Clonality)	Polyclonal
剂型(Form)	Liquid
保存条件(Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液(Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化 (Purification)	Affinity purification

产品应用(Application)

稀释比(Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:10000
分子量(Molecular Weight)	80kDa

抗原信息(Antigen Information)

基因名(Gene Name)	GNE
别名(Alternative Names)	GNE; GLCNE; Bifunctional UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase; UDP-GlcNAc-2-epimerase/ManAc kinase
基因 ID(Gene ID)	10020.0
SwissProt ID	Q9Y223
免疫原(Immunogen)	The antiserum was produced against synthesized peptide derived from human GNE. AA range:592-641

研究背景 (Background)

The protein encoded by this gene is a bifunctional enzyme that initiates and regulates the biosynthesis of N-acetylneuraminic

acid (NeuAc), a precursor of sialic acids. It is a rate-limiting enzyme in the sialic acid biosynthetic pathway. Sialic acid modification of cell surface molecules is crucial for their function in many biologic processes, including cell adhesion and signal transduction. Differential sialylation of cell surface molecules is also implicated in the tumorigenicity and metastatic behavior of malignant cells. Mutations in this gene are associated with sialuria, autosomal recessive inclusion body myopathy, and Nonaka myopathy. Alternative splicing of this gene results in transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008],catalytic activity:ATP + N-acetyl-D-mannosamine = ADP + N-acetyl-D-mannosamine 6-phosphate.,catalytic activity:UDP-N-acetyl-D-glucosamine = UDP-N-acetyl-D-mannosamine.,disease:Defects in GNE are a cause of sialuria [MIM:269921]; also known as sialuria French type. In sialuria, free sialic acid accumulates in the cytoplasm and gram quantities of neuraminic acid are secreted in the urine. The metabolic defect involves lack of feedback inhibition of UDP-GlcNAc 2-epimerase by CMP-Neu5Ac, resulting in constitutive overproduction of free Neu5Ac. Clinical features include variable degrees of developmental delay, coarse facial features and hepatomegaly. Sialuria inheritance is autosomal dominant.,disease:Defects in GNE are the cause of inclusion body myopathy type 2 (IBM2) [MIM:600737]. Hereditary inclusion body myopathies are a group of neuromuscular disorders characterized by adult onset, slowly progressive distal and proximal weakness and a typical muscle pathology including rimmed vacuoles and filamentous inclusions. IBM2 is an autosomal recessive disorder affecting mainly leg muscles, but with an unusual distribution that spares the quadriceps as also observed in Nonaka myopathy.,disease:Defects in GNE are the cause of Nonaka myopathy (NM) [MIM:605820]; also known as distal myopathy with rimmed vacuoles (DMRV). NM is an autosomal recessive muscular disorder, allelic to inclusion body myopathy 2. It is characterized by weakness of the anterior compartment of the lower limbs with onset in early adulthood, and sparing of the quadriceps muscles. As the inclusion body myopathy, NM is histologically characterized by the presence of numerous rimmed vacuoles without inflammatory changes in muscle specimens.,enzyme regulation:Allosterically regulated (Probable); feedback inhibited by cytidine monophosphate-N-acetylneuraminic acid (CMP-Neu5Ac), the end product of neuraminic acid biosynthesis. Activity is dependent on oligomerization. The monomer is inactive, whereas the dimer catalyzes only the phosphorylation of N-acetylmannosamine; the hexamer is fully active for both enzyme activities (By similarity). Up-regulated after PKC-dependent phosphorylation.,function:Regulates and initiates biosynthesis of N-acetylneuraminic acid (NeuAc), a precursor of sialic acids. Plays an essential role in early development (By similarity). Required for normal sialylation in hematopoietic cells. Sialylation is implicated in cell adhesion, signal transduction, tumorigenicity and metastatic behavior of malignant cells.,pathway:Amino-sugar metabolism; N-acetylneuraminic acid biosynthesis.,PTM:Phosphorylated by PKC.,similarity:In the C-terminal section; belongs to the ROK (nagC/xyIR) family.,similarity:In the N-terminal section; belongs to the UDP-N-acetylglucosamine 2-epimerase family.,subunit:Homodimer and homohexamer.,tissue specificity:Highest expression in liver and placenta. Also found in heart, brain, lung, kidney, skeletal muscle and pancreas.,


研究领域 (Research Area)

Amino sugar and nucleotide sugar metabolism;

注意事项 (Note)

For research use only.

图片 (Image Data)



Western Blot analysis of 3T3 cells using GLCNE Polyclonal Antibody diluted at 1 : 500

EnkiLife 优势产品推荐

产品类别	产品货号	产品名称
WB 解决方案	RA10020	2h 极速 WB 即用型全流程试剂盒
	RA10021	4h 快速 WB 即用型全流程试剂盒
	RA10042	5h 畅享版 WB 全流程试剂盒
	RA10037	校准级彩色预染蛋白 Marker (8-180kDa)
	RA10038	校准级彩色预染蛋白 Marker (10-250kDa)
	RA10039	校准级高分子彩色预染蛋白 Marker (25-400kDa)
TSA 多重荧光染色试剂盒	RA10008	TSA 双标三色多重荧光染色试剂盒 (mIHC)
	RA10009	TSA 三标四色多重荧光染色试剂盒 (mIHC)
	RA10010	TSA 四标五色多重荧光染色试剂盒 (mIHC)
	RA10011	TSA 五标六色多重荧光染色试剂盒 (mIHC)
	RA10012	TSA 六标七色多重荧光染色试剂盒 (mIHC)
IHC 检测试剂盒	RA10006	HRP Anti-Mouse/Rabbit IHC Detection System
	RA10007	Polymer-HRP Anti-Mouse/Rabbit IHC Detection System
抗体标记试剂盒	RE80004p	辣根过氧化物酶(HRP)抗体标记试剂盒
	RE80002q	Sulfo-NHS-生物素标记试剂盒
	RE80007p	Cy3 荧光素标记试剂盒
	RE80011p	Fluor488 荧光素标记试剂盒
	RE80017p	Fluor750 荧光素标记试剂盒
	RE80005p	藻红蛋白(R-PE) 抗体快速标记试剂盒
	RE80040	PE-Cy7 串联染料抗体快速标记试剂盒
稳转细胞系构建服务 (免费赠送全膜 WB 验证)	TS-0001	过表达稳转细胞系构建
	TS-0002	敲低稳转细胞系构建
	TS-0003	敲除细胞系构建

限于篇幅，更多产品详见官网：<https://www.enkilife.cn>