

产品名称: GPR37 Rabbit Polyclonal Antibody
产品货号: AP Rab11678



产品概述 (Summary)

产品名称 (Production Name)	GPR37 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	GPR37 GPR37; Probable G-protein coupled receptor 37; Endothelin B receptor-like protein 1; ETBR-LP-1; Parkin-associated endothelin receptor-like receptor; PAELR
别名 (Alternative Names)	
基因 ID (Gene ID)	2861.0
蛋白 ID (SwissProt ID)	O15354.The antiserum was produced against synthesized peptide derived from human GPR37. AA range:211-260

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	68kDa

产品名称: GPR37 Rabbit Polyclonal Antibody
产品货号: AP Rab11678




研究背景 (Background)

This gene is a member of the G protein-coupled receptor family. The encoded protein contains seven transmembrane domains and is found in cell and endoplasmic reticulum membranes. G protein-coupled receptors are involved in translating outside signals into G protein mediated intracellular effects. This gene product interacts with Parkin and is involved in juvenile Parkinson disease. [provided by RefSeq, Oct 2012],function:Orphan receptor. May have a unique functional role in the central nervous system.,PTM:Ubiquitinated by PARK2 in the presence of UBE2E1 and UBE2L3 in the endoplasmic reticulum. The unfolded form is specifically ubiquitinated by SYVN1, which promotes its proteasomal degradation and prevents neuronal cell death.,similarity:Belongs to the G-protein coupled receptor 1 family.,subunit:Forms a complex with PARK2, STUB1 and HSP70. The amount of STUB1 in the complex increases during ER stress. STUB1 promotes the dissociation of HSP70 from PARK2, thus facilitating PARK2-mediated GPR37 ubiquitination. Interacts with PACRG.,tissue specificity:Expressed in brain and spinal cord, and at lower levels in testis, placenta and liver, but no detectable expression observed in any other tissue. When overexpressed in cells, tends to become insoluble and unfolded. Accumulation of the unfolded protein may lead to dopaminergic neuronal death in juvenile Parkinson disease (PDJ),

研究领域 (Research Area)


Parkinson's disease;

图片 (Image Data)



Western blot analysis of GPR37 Antibody. The lane on the right is blocked with the GPR37 peptide.

产品名称: GPR37 Rabbit Polyclonal Antibody
产品货号: APRab11678



Western blot analysis of the lysates from HT-29 cells using GPR37 antibody.

注意事项 (Note)

For research use only .