产品货号: APRab11716



## 产品概述 (Summary)

产品名称 (Production Name) GPRC6A Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application)WB,ICC/IF,ELISA种属反应性 (Reactivity)Human,Rat,Mouse

## 产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

# 免疫原信息 (Immunogen)

基因名 (Gene Name) GPRC6A

GPRC6A; G-protein coupled receptor family C group 6 member A; hGPRC6A; 别名 (Alternative Names)

G-protein coupled receptor GPCR33; hGPCR33

基因 ID (Gene ID) 222545.0

Q5T6X5.The antiserum was produced against synthesized peptide derived 蛋白ID (SwissProt ID)

from human GPRC6A. AA range:471-520

## 产品应用 (Application)

**稀释比 (Dilution Ratio)** WB 1:500-1:2000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:10000

蛋白分子量 (Molecular Weight) 105kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品货号: APRab11716




#### 研究背景 (Background)

Members of family C of the G protein-coupled receptor (GPCR) superfamily, such as GPRC6A, are characterized by an evolutionarily conserved amino acid-sensing motif linked to an intramembranous 7-transmembrane loop region. Several members of GPCR family C, including GPRC6A, also have a long N-terminal domain (summary by Pi et al., 2005 [PubMed 16199532]). [supplied by OMIM, Nov 2010], function: Receptor that is activated by both amino acids and extracellular concentration of calcium ions. The activity of this receptor is mediated by a G-protein that activates a phosphatidylinositol-calcium second messenger system. Senses changes in the extracellular concentration of calcium ions, suggesting that it may mediate extracellular calcium-sensing responses in osteoblasts. Osteocalin, stimulates its activity in presence of calcium. Has a lower affinity for calcium than CASR. Also acts as a receptor for amino acids, with a preference for basic amino acids such as L-Lys, L-Arg and L-ornithine. Its affinity for amino acids suggests that it may act as a regulatory component of the urea cycle., PTM:N-glycosylated., similarity:Belongs to the G-protein coupled receptor 3 family., subunit:Homodimer; disulfide-linked., tissue specificity:Widely expressed. Expressed at high level in brain, skeletal muscle, testis, bone, calvaria, osteoblasts and leukocytes. Expressed at intermediate level in liver, heart, kidney and spleen. Expressed at low level in lung, pancreas, placenta and ovary. Not expressed in thymus, prostate, small intestine, tongue and colon. In kidney, isoform 1 and isoform 2 are expressed at the same level, while in other tissues, isoform 2 is expressed at low level.

## 研究领域 (Research Area)

# 图片 (Image Data)




Immunofluorescence analysis of MCF7 cells, using GPRC6A Antibody. The picture on the right is blocked with the synthesized peptide.


Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品货号: APRab11716






Western blot analysis of lysates from Jurkat cells, using GPRC6A Antibody. The lane on the right is blocked with the synthesized peptide.




Western blot analysis of the lysates from HeLa cells using GPRC6A antibody.



Western Blot analysis of Jurkat cells using GPRC6A Polyclonal Antibody diluted at 1: 1000

产品货号: APRab11716





Western Blot analysis of various cells using Antibody diluted at 1:1000. Secondary antibody was diluted at 1:20000

## 注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838