

产品名称: GRIP-1 Rabbit Polyclonal Antibody

产品货号: APRab11770

产品概述 (Summary)

描述(Description)	Rabbit polyclonal Antibody
宿主(Host)	Rabbit
应用(Application)	IHC, ICC/IF, ELISA
种属反应(Reactivity)	Human, Rat, Mouse
偶联物(Conjugation)	Unconjugated
修饰(Modification)	Unmodified
同种型(Isotype)	IgG
克隆(Clonality)	Polyclonal
剂型(Form)	Liquid
保存条件(Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液(Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化 (Purification)	Affinity purification

产品应用(Application)

稀释比(Dilution Ratio) IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:20000-1:40000

分子量(Molecular Weight)

抗原信息(Antigen Information)

基因名(Gene Name)	NCOA2
别名(Alternative Names)	NCOA2; BHLHE75; TIF2; Nuclear receptor coactivator 2; NCoA-2; Class E basic helix-loop-helix protein 75; bHLHe75; Transcriptional intermediary factor 2; hTIF2
基因 ID(Gene ID)	10499.0
SwissProt ID	Q15596
免疫原(Immunogen)	The antiserum was produced against synthesized peptide derived from human NCoA2. AA range:702-751

研究背景 (Background)

The protein encoded by this gene functions as a transcriptional coactivator for nuclear hormone receptors, including steroid,


thyroid, retinoid, and vitamin D receptors. The encoded protein acts as an intermediary factor for the ligand-dependent activity of these nuclear receptors, which regulate their target genes upon binding of cognate response elements. This gene has been found to be involved in translocations that result in fusions with other genes in various cancers, including the lysine acetyltransferase 6A (KAT6A) gene in acute myeloid leukemia, the ETS variant 6 (ETV6) gene in acute lymphoblastic leukemia, and the related family bHLH transcription factor with YRPW motif 1 (HEY1) gene in mesenchymal chondrosarcoma. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016],disease:Chromosomal aberrations involving NCOA2 may be a cause of acute myeloid leukemias. Inversion inv(8)(p11;q13) generates the MYST3-NCOA2 oncogene, which consists of the N-terminus part of MYST3/MOZ and the C-terminus part of NCOA2/TIF2. MYST3-NCOA2 binds to CREBBP and disrupts its function in transcription activation.,domain:Contains 2 C-terminal transcription activation domains (AD1 and AD2) that can function independently.,domain:Contains four Leu-Xaa-Xaa-Leu-Leu (LXXLL) motifs. The LXXLL motifs are essential for the association with nuclear receptors and are, at least in part, functionally redundant.,domain:The LLXXLXXLL motif is involved in transcriptional coactivation and CREBBP/CBP binding.,function:Transcriptional coactivator for steroid receptors and nuclear receptors. Coactivator of the steroid binding domain (AF-2) but not of the modulating N-terminal domain (AF-1). Required with NCOA1 to control energy balance between white and brown adipose tissues.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the SRC/p160 nuclear receptor coactivator family.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,similarity:Contains 1 PAS (PER-ARNT-SIM) domain.,subunit:Present in a complex containing CARM1 and EP300/P300, and interacts with CARM1 and NR3C2 (By similarity). Present in a complex containing NCOA3, IKKA, IKKB, IKBKG and CREBBP. Interacts (via C-terminus) with CREBBP. Interacts with HIF1A, NCOA1, APEX and NR3C1. Interacts with CASP8AP2 and TTL5/STAMP. Interacts with ESR1, RARA and RXRA.,

研究领域 (Research Area)

注意事项 (Note)

For research use only.

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200 (4° overnight) . 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min) .

EnkiLife 优势产品推荐

产品类别	产品货号	产品名称
WB 解决方案	RA10020	2h 极速 WB 即用型全流程试剂盒
	RA10021	4h 快速 WB 即用型全流程试剂盒
	RA10042	5h 畅享版 WB 全流程试剂盒
	RA10037	校准级彩色预染蛋白 Marker (8-180kDa)
	RA10038	校准级彩色预染蛋白 Marker (10-250kDa)
	RA10039	校准级高分子彩色预染蛋白 Marker (25-400kDa)
TSA 多重荧光染色试剂盒	RA10008	TSA 双标三色多重荧光染色试剂盒 (mIHC)
	RA10009	TSA 三标四色多重荧光染色试剂盒 (mIHC)
	RA10010	TSA 四标五色多重荧光染色试剂盒 (mIHC)
	RA10011	TSA 五标六色多重荧光染色试剂盒 (mIHC)
	RA10012	TSA 六标七色多重荧光染色试剂盒 (mIHC)
IHC 检测试剂盒	RA10006	HRP Anti-Mouse/Rabbit IHC Detection System
	RA10007	Polymer-HRP Anti-Mouse/Rabbit IHC Detection System
抗体标记试剂盒	RE80004p	辣根过氧化物酶(HRP)抗体标记试剂盒
	RE80002q	Sulfo-NHS-生物素标记试剂盒
	RE80007p	Cy3 荧光素标记试剂盒
	RE80011p	Fluor488 荧光素标记试剂盒
	RE80017p	Fluor750 荧光素标记试剂盒
	RE80005p	藻红蛋白(R-PE) 抗体快速标记试剂盒
	RE80040	PE-Cy7 串联染料抗体快速标记试剂盒
稳转细胞系构建服务 (免费赠送全膜 WB 验证)	TS-0001	过表达稳转细胞系构建
	TS-0002	敲低稳转细胞系构建
	TS-0003	敲除细胞系构建

限于篇幅，更多产品详见官网：<https://www.enkilife.cn>