

产品名称: GRK 2 Rabbit Polyclonal Antibody
产品货号: APRab11773



产品概述 (Summary)

产品名称 (Production Name)	GRK 2 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ADRBK1
别名 (Alternative Names)	ADRBK1; BARK; BARK1; GRK2; Beta-adrenergic receptor kinase 1; Beta-ARK-1; G-protein coupled receptor kinase 2
基因 ID (Gene ID)	156.0
蛋白 ID (SwissProt ID)	P25098.The antiserum was produced against synthesized peptide derived from human ARBK1. AA range:601-650

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	80kDa

产品名称: GRK 2 Rabbit Polyclonal Antibody
产品货号: AP Rab11773





研究背景 (Background)

The product of this gene phosphorylates the beta-2-adrenergic receptor and appears to mediate agonist-specific desensitization observed at high agonist concentrations. This protein is an ubiquitous cytosolic enzyme that specifically phosphorylates the activated form of the beta-adrenergic and related G-protein-coupled receptors. Abnormal coupling of beta-adrenergic receptor to G protein is involved in the pathogenesis of the failing heart. [provided by RefSeq, Jul 2008], catalytic activity: ATP + [beta-adrenergic receptor] = ADP + [beta-adrenergic receptor] phosphate., catalytic activity: ATP + a protein = ADP + a phosphoprotein., function: Specifically phosphorylates the agonist-occupied form of the beta-adrenergic and closely related receptors, probably inducing a desensitization of them., online information: Beta adrenergic receptor kinase entry, similarity: Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. GPRK subfamily., similarity: Contains 1 AGC-kinase C-terminal domain., similarity: Contains 1 PH domain., similarity: Contains 1 protein kinase domain., similarity: Contains 1 RGS domain., subunit: Interacts with GIT1 (By similarity). Interacts with, and phosphorylates chemokine-stimulated CCR5., tissue specificity: Expressed in peripheral blood leukocytes.,

研究领域 (Research Area)

Chemokine; Endocytosis;


图片 (Image Data)




产品名称: GRK 2 Rabbit Polyclonal Antibody
产品货号: AP Rab11773




Western Blot analysis of various cells using GRK 2 Polyclonal Antibody



Western Blot analysis of HuvEc cells using GRK 2 Polyclonal Antibody




Immunohistochemical analysis of paraffin-embedded Human lung cancer. Antibody was diluted at 1:100 (4°, overnight). High-pressure and temperature Tris-EDTA, pH 8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.



Immunohistochemical analysis of paraffin-embedded Human Lymph gland. 1, Antibody was diluted at 1:100 (4°, overnight). 2, High-pressure and temperature EDTA, pH 8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 30min).

产品名称: GRK 2 Rabbit Polyclonal Antibody
产品货号: APRab11773



Immunohistochemical analysis of paraffin-embedded Human Lymph gland. 1, Antibody was diluted at 1:100 (4°,overnight) .
2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at
1:200 (room temperature, 30min) .

注意事项 (Note)

For research use only .