

产品名称: GRO α Rabbit Polyclonal Antibody
产品货号: APRab11788



产品概述 (Summary)

产品名称 (Production Name)	GRO α Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CXCL1 CXCL1; GRO; GRO1; GROA; MGSA; SCYB1; Growth-regulated alpha protein; C-X-C motif chemokine 1; GRO-alpha(1-73); Melanoma growth stimulatory activity; MGSA; Neutrophil-activating protein 3; NAP-3
别名 (Alternative Names)	
基因 ID (Gene ID)	2919.0
蛋白 ID (SwissProt ID)	P09341. The antiserum was produced against synthesized peptide derived from human GROalpha. AA range: 39-88

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	

产品名称: GRO α Rabbit Polyclonal Antibody
产品货号: AP Rab11788




研究背景 (Background)

This antimicrobial gene encodes a member of the CXC subfamily of chemokines. The encoded protein is a secreted growth factor that signals through the G-protein coupled receptor, CXC receptor 2. This protein plays a role in inflammation and as a chemoattractant for neutrophils. Aberrant expression of this protein is associated with the growth and progression of certain tumors. A naturally occurring processed form of this protein has increased chemotactic activity. Alternate splicing results in coding and non-coding variants of this gene. A pseudogene of this gene is found on chromosome 4. [provided by RefSeq, Sep 2014],function:Has chemotactic activity for neutrophils. May play a role in inflammation and exerts its effects on endothelial cells in an autocrine fashion. In vitro, the processed forms GRO-alpha(4-73), GRO-alpha(5-73) and GRO-alpha(6-73) show a 30-fold higher chemotactic activity.,online information: CXCL1 entry,PTM:N-terminal processed forms GRO-alpha(4-73), GRO-alpha(5-73) and GRO-alpha(6-73) are produced by proteolytic cleavage after secretion from peripheral blood monocytes.,similarity:Belongs to the intercrine alpha (chemokine CxC) family.,

研究领域 (Research Area)

Cytokine-cytokine receptor interaction;Chemokine;NOD-like receptor;Epithelial cell signaling in Helicobacter pylori infection;

图片 (Image Data)



Immunohistochemistry analysis of GRO α antibody in paraffin-embedded human brain tissue.

注意事项 (Note)

For research use only .