产品名称: hCAP-G Rabbit Polyclonal Antibody

产品货号: APRab11920



产品概述 (Summary)

产品名称 (Production Name) hCAP-G Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit 应用 (Application) WB,ELISA

种属反应性 (Reactivity) Human, Rat, Mouse

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) NCAPG

NCAPG; CAPG; NYMEL3; Condensin complex subunit 3; Chromosome-

别名 (Alternative Names) associated protein G; Condensin subunit CAP-G; hCAP-G; Melanoma antigen

NY-MEL-3; Non-SMC condensin I complex subunit G; XCAP-G homolog

基**因 ID (Gene ID)** 64151.0

Q9BPX3.The antiserum was produced against synthesized peptide derived 蛋白ID (SwissProt ID)

from human NCAPG. AA range:951-1000

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,ELISA 1:20000-1:40000

蛋白分子量 (Molecular Weight) 115kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: hCAP-G Rabbit Polyclonal Antibody

产品货号: APRab11920




研究背景 (Background)

This gene encodes a subunit of the condensin complex, which is responsible for the condensation and stabilization of chromosomes during mitosis and meiosis. Phosphorylation of the encoded protein activates the condensin complex. There are pseudogenes for this gene on chromosomes 8 and 15. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2012], function: Regulatory subunit of the condensin complex, a complex required for conversion of interphase chromatin into mitotic-like condense chromosomes. The condensin complex probably introduces positive supercoils into relaxed DNA in the presence of type I topoisomerases and converts nicked DNA into positive knotted forms in the presence of type II topoisomerases, miscellaneous: Overexpressed in some cancer lines and some tumor cells, PTM: Phosphorylated by CDC2. Its phosphorylation, as well as that of NCAPD2 and NCAPH subunits, activates the condensin complex and is required for chromosome condensation, sequence caution: Wrong choice of frame., similarity: Belongs to the CND3 (condensin subunit 3) family., similarity: Contains 10 HEAT repeats., subcellular location:In interphase cells, the majority of the condensin complex is found in the cytoplasm, while a minority of the complex is associated with chromatin. A subpopulation of the complex however remains associated with chromosome foci in interphase cells. During mitosis, most of the condensin complex is associated with the chromatin. At the onset of prophase, the regulatory subunits of the complex are phosphorylated by CDC2, leading to condensin's association with chromosome arms and to chromosome condensation. Dissociation from chromosomes is observed in late telophase., subunit: Component of the condensin complex, which contains the SMC2 and SMC4 heterodimer, and three non SMC subunits that probably regulate the complex: NCAPH/BRRN1, NCAPD2/CAPD2 and NCAPG, tissue specificity: Highly expressed in testis.,

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of NCAPG Antibody. The lane on the right is blocked with the NCAPG peptide.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品名称: hCAP-G Rabbit Polyclonal Antibody

产品货号: APRab11920



注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838