

产品名称: hCAP-H Rabbit Polyclonal Antibody
产品货号: AP Rab11922



产品概述 (Summary)

产品名称 (Production Name)	hCAP-H Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	NCAPH NCAPH; BRRN; BRRN1; CAPH; KIAA0074; Condensin complex subunit 2; Barren
别名 (Alternative Names)	homolog protein 1; Chromosome-associated protein H; hCAP-H; Non-SMC condensin I complex subunit H; XCAP-H homolog
基因 ID (Gene ID)	23397.0
蛋白 ID (SwissProt ID)	Q15003.The antiserum was produced against synthesized peptide derived from human NCAPH. AA range:441-490

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	83kDa

产品名称: hCAP-H Rabbit Polyclonal Antibody
产品货号: AP Rab11922




研究背景 (Background)

This gene encodes a member of the barr gene family and a regulatory subunit of the condensin complex. This complex is required for the conversion of interphase chromatin into condensed chromosomes. The protein encoded by this gene is associated with mitotic chromosomes, except during the early phase of chromosome condensation. During interphase, the protein has a distinct punctate nucleolar localization. Alternatively spliced transcript variants encoding different proteins have been described. [provided by RefSeq, Jul 2013],function:Regulatory subunit of the condensin complex, a complex required for conversion of interphase chromatin into mitotic-like condense chromosomes. The condensin complex probably introduces positive supercoils into relaxed DNA in the presence of type I topoisomerases and converts nicked DNA into positive knotted forms in the presence of type II topoisomerases.,PTM:Phosphorylated by CDC2. Its phosphorylation, as well as that of NCAPD2 and NCAPG subunits, activates the condensin complex and is required for chromosome condensation.,similarity:Belongs to the CND2 (condensin subunit 2) family.,subcellular location:In interphase cells, the majority of the condensin complex is found in the cytoplasm, while a minority of the complex is associated with chromatin. A subpopulation of the complex however remains associated with chromosome foci in interphase cells. During mitosis, most of the condensin complex is associated with the chromatin. At the onset of prophase, the regulatory subunits of the complex are phosphorylated by CDC2, leading to condensin's association with chromosome arms and to chromosome condensation. Dissociation from chromosomes is observed in late telophase.,subunit:Component of the condensin complex, which contains the SMC2 and SMC4 heterodimer, and three non SMC subunits that probably regulate the complex: NCAPH/BRRN1, NCAPD2/CAPD2 and NCAPG.,tissue specificity:Widely expressed at low level. Expressed in proliferating cells.,


研究领域 (Research Area)

图片 (Image Data)




Western blot analysis of lysates from COLO cells, using NCAPH Antibody. The lane on the right is blocked with the synthesized peptide.

产品名称: hCAP-H Rabbit Polyclonal Antibody
产品货号: APRab11922



Western blot analysis of the lysates from COLO205 cells using NCAPH antibody.



Western Blot analysis of COLO cells using hCAP-H Polyclonal Antibody

注意事项 (Note)

For research use only .