

产品名称: HCF1 Rabbit Polyclonal Antibody

产品货号: APRab11926

产品概述 (Summary)

描述(Description)	Rabbit polyclonal Antibody
宿主(Host)	Rabbit
应用(Application)	WB,IHC,ICC/IF,ELISA
种属反应(Reactivity)	Human,Mouse
偶联物(Conjugation)	Unconjugated
修饰(Modification)	Unmodified
同种型(Isotype)	IgG
克隆(Clonality)	Polyclonal
剂型(Form)	Liquid
保存条件(Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液(Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化 (Purification)	Affinity purification

产品应用(Application)

稀释比(Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
分子量(Molecular Weight)	208kDa

抗原信息(Antigen Information)

基因名(Gene Name)	HCFC1
别名(Alternative Names)	HCFC1; HCF1; HFC1; Host cell factor 1; HCF; HCF-1; C1 factor; CFF; VCAF; VP16 accessory protein
基因 ID(Gene ID)	3054.0
SwissProt ID	P51610
免疫原(Immunogen)	The antiserum was produced against synthesized peptide derived from human HCFC1. AA range:131-180

研究背景 (Background)

This gene is a member of the host cell factor family and encodes a protein with five Kelch repeats, a fibronectin-like motif, and


six HCF repeats, each of which contains a highly specific cleavage signal. This nuclear coactivator is proteolytically cleaved at one of the six possible sites, resulting in the creation of an N-terminal chain and the corresponding C-terminal chain. The final form of this protein consists of noncovalently bound N- and C-terminal chains. The protein is involved in control of the cell cycle and transcriptional regulation during herpes simplex virus infection. Alternatively spliced variants which encode different protein isoforms have been described; however, not all variants have been fully characterized. [provided by RefSeq, Jul 2008],domain:The HCF repeat is a highly specific proteolytic cleavage signal.,domain:The kelch repeats fold into a 6-bladed kelch beta-propeller called the beta-propeller domain which mediates interaction with HCFC1R1.,function:Involved in control of the cell cycle. Upon lytic infection of permissive cells, the HSV transactivator protein VP16 associates with HCFC1. Binding to HCFC1 activates VP16 for association with the octamer motif-binding protein POU2F1, to form a multiprotein-DNA complex responsible for activating transcription of the HSV immediate early genes. Also antagonizes transactivation by ZBTB17 and GABP2; represses ZBTB17 activation of the p15(INK4b) promoter and inhibits its ability to recruit p300. Coactivator for EGR2 and GABP2. Tethers the chromatin modifying Set1/Ash2 histone H3-K4 methyltransferase (HMT) and Sin3 histone deacetylase (HDAC) complexes (involved in the activation and repression of transcription, respectively) together.,PTM:O-glycosylated.,PTM:Proteolytically cleaved at one or several PPCE--THET sites within the HCF repeats. Further cleavage of the primary N- and C-terminal chains results in a 'trimming' and accumulation of the smaller chains.,similarity:Contains 5 Kelch repeats.,subcellular location:HCFC1R1 modulates its subcellular localization and overexpression of HCFC1R1 leads to accumulation of HCFC1 in the cytoplasm.,subunit:Composed predominantly of six polypeptides ranging from 110 to 150 kDa and a minor 300 kDa polypeptide. The majority of N-and C-terminal cleavage products remain tightly, albeit non-covalently, associated. Interacts with POU2F1, VP16, CREB3, ZBTB17, EGR2, E2F4, CREBZF, SP1, GABP2, Sin3 HDAC complex (SIN3A, HDAC1, HDAC2, SDS3), Set1/Ash2 HMT complex (SET1, ASH2, WDR5), SAP30, SIN3B, OGT1 and FHL2. Component of the MLL complex, at least composed of MLL, ASH2L, RBBP5, DPY30, WDR5, MEN1, HCFC1 and HCFC2. Interacts with HCFC1R1 and THAP11.,tissue specificity:Highly expressed in fetal tissues and the adult kidney. Present in all tissues tested.,

研究领域 (Research Area)

注意事项 (Note)

For research use only.

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using HCFC1 Antibody. The picture on the right is blocked with the synthesized peptide.

EnkiLife 优势产品推荐

产品类别	产品货号	产品名称
WB 解决方案	RA10020	2h 极速 WB 即用型全流程试剂盒
	RA10021	4h 快速 WB 即用型全流程试剂盒
	RA10042	5h 畅享版 WB 全流程试剂盒
	RA10037	校准级彩色预染蛋白 Marker (8-180kDa)
	RA10038	校准级彩色预染蛋白 Marker (10-250kDa)
	RA10039	校准级高分子彩色预染蛋白 Marker (25-400kDa)
TSA 多重荧光染色试剂盒	RA10008	TSA 双标三色多重荧光染色试剂盒 (mIHC)
	RA10009	TSA 三标四色多重荧光染色试剂盒 (mIHC)
	RA10010	TSA 四标五色多重荧光染色试剂盒 (mIHC)
	RA10011	TSA 五标六色多重荧光染色试剂盒 (mIHC)
	RA10012	TSA 六标七色多重荧光染色试剂盒 (mIHC)
IHC 检测试剂盒	RA10006	HRP Anti-Mouse/Rabbit IHC Detection System
	RA10007	Polymer-HRP Anti-Mouse/Rabbit IHC Detection System
抗体标记试剂盒	RE80004p	辣根过氧化物酶(HRP)抗体标记试剂盒
	RE80002q	Sulfo-NHS-生物素标记试剂盒
	RE80007p	Cy3 荧光素标记试剂盒
	RE80011p	Fluor488 荧光素标记试剂盒
	RE80017p	Fluor750 荧光素标记试剂盒
	RE80005p	藻红蛋白(R-PE) 抗体快速标记试剂盒
	RE80040	PE-Cy7 串联染料抗体快速标记试剂盒
稳转细胞系构建服务 (免费赠送全膜 WB 验证)	TS-0001	过表达稳转细胞系构建
	TS-0002	敲低稳转细胞系构建
	TS-0003	敲除细胞系构建

限于篇幅，更多产品详见官网：<https://www.enkilife.cn>