

产品名称: IgG1 Rabbit Polyclonal Antibody
产品货号: APRab12443



产品概述 (Summary)

产品名称 (Production Name)	IgG1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	IGHG1
别名 (Alternative Names)	IGHG1; Ig gamma-1 chain C region
基因 ID (Gene ID)	3500.0
蛋白 ID (SwissProt ID)	P01857.The antiserum was produced against synthesized peptide derived from human IgG1. AA range:196-245

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	41kDa

研究背景 (Background)

产品名称: IgG1 Rabbit Polyclonal Antibody
产品货号: APRab12443




disease:Chromosomal aberrations involving IGHG1 may be a cause of multiple myeloma [MIM:254500]. Translocation t(11;14)(q13;q32) with CCND1; translocation t(4;14)(p16.3;q32.3) with FGFR3; translocation t(6;14)(p25;q32) with IRF4.,miscellaneous:Disease protein OMM may represent an allelic form or another gamma chain subclass.,miscellaneous:Disease protein WIS is lacking most of the V region and all of the CH1 region.,miscellaneous:Disease protein ZUC lack most of the V region, all of the CH1 region, and part of the hinge compared with normal gamma-3 heavy chains.,miscellaneous:EU also differs in the amidation states of residues 155, 166, 177, 195, 198, 269, and 272 and in the order of residues 268-272.,miscellaneous:KOL also differs in the amidation states of residues 198, 267 and 272.,miscellaneous:Nie also differs in the amidation states of 35, 116, 198, 269 and 272.,miscellaneous:Nie has the G1M(17) allotypic marker, 97-K, and the G1M(1) markers, 239-D and 241-L. KOL and EU sequences have the G1M(3) marker and the G1M (non-1) markers.,miscellaneous:The hinge region in gamma-3 chains is about four times as long as in other gamma chains and contains three identical 15-residue segments preceded by a similar 17-residue segment (12-28).,online information:IGHM mutation db,polymorphism:All 4 combinations of the S/G and V/G polymorphisms at positions 191 and 216 have been observed in human mu chains.,subcellular location:During differentiation, B-lymphocytes switch from expression of membrane-bound IgM to secretion of IgM.,subunit:Dimer linked by 12 disulfide bonds; it has an extra interchain disulfide bond at position 7 in addition to the 11 normally present in the hinge region.,disease:Chromosomal aberrations involving IGHG1 may be a cause of multiple myeloma [MIM:254500]. Translocation t(11;14)(q13;q32) with CCND1; translocation t(4;14)(p16.3;q32.3) with FGFR3; translocation t(6;14)(p25;q32) with IRF4.,miscellaneous:Disease protein OMM may represent an allelic form or another gamma chain subclass.,miscellaneous:Disease protein WIS is lacking most of the V region and all of the CH1 region.,miscellaneous:Disease protein ZUC lack most of the V region, all of the CH1 region, and part of the hinge compared with normal gamma-3 heavy chains.,miscellaneous:EU also differs in the amidation states of residues 155, 166, 177, 195, 198, 269, and 272 and in the order of residues 268-272.,miscellaneous:KOL also differs in the amidation states of residues 198, 267 and 272.,miscellaneous:Nie also differs in the amidation states of 35, 116, 198, 269 and 272.,miscellaneous:Nie has the G1M(17) allotypic marker, 97-K, and the G1M(1) markers, 239-D and 241-L. KOL and EU sequences have the G1M(3) marker and the G1M (non-1) markers.,miscellaneous:The hinge region in gamma-3 chains is about four times as long as in other gamma chains and contains three identical 15-residue segments preceded by a similar 17-residue segment (12-28).,online information:IGHM mutation db,polymorphism:All 4 combinations of the S/G and V/G polymorphisms at positions 191 and 216 have been observed in human mu chains.,subcellular location:During differentiation, B-lymphocytes switch from expression of membrane-bound IgM to secretion of IgM.,subunit:Dimer linked by 12 disulfide bonds; it has an extra interchain disulfide bond at position 7 in addition to the 11 normally present in the hinge region.,


研究领域 (Research Area)

图片 (Image Data)


产品名称: IgG1 Rabbit Polyclonal Antibody
产品货号: APRab12443



Immunohistochemistry analysis of IgG1 antibody in paraffin-embedded human lung carcinoma tissue.



Western blot analysis of lysate from LOVO cells, using IgG1 antibody.



Western Blot analysis of colo cells using IgG1 Polyclonal Antibody diluted at 1 : 500

注意事项 (Note)

For research use only .