

产品名称: IL-15R α Rabbit Polyclonal Antibody
产品货号: APRab12502



产品概述 (Summary)

产品名称 (Production Name)	IL-15R α Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	IL15RA
别名 (Alternative Names)	IL15RA; Interleukin-15 receptor subunit alpha; IL-15 receptor subunit alpha; IL-15R-alpha; IL-15RA; CD antigen CD215
基因 ID (Gene ID)	3601.0
蛋白 ID (SwissProt ID)	Q13261.The antiserum was produced against synthesized peptide derived from human IL15RA. AA range:99-148

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	32kDa

产品名称: IL-15R α Rabbit Polyclonal Antibody
产品货号: AP Rab12502




研究背景 (Background)

This gene encodes a cytokine receptor that specifically binds interleukin 15 (IL15) with high affinity. The receptors of IL15 and IL2 share two subunits, IL2R beta and IL2R gamma. This forms the basis of many overlapping biological activities of IL15 and IL2. The protein encoded by this gene is structurally related to IL2R alpha, an additional IL2-specific alpha subunit necessary for high affinity IL2 binding. Unlike IL2RA, IL15RA is capable of binding IL15 with high affinity independent of other subunits, which suggests distinct roles between IL15 and IL2. This receptor is reported to enhance cell proliferation and expression of apoptosis inhibitor BCL2L1/BCL2-XL and BCL2. Multiple alternatively spliced transcript variants of this gene have been reported.[provided by RefSeq, Apr 2010],function:Receptor for interleukin-15. Expression of different isoforms may alter or interfere with signal transduction. Isoform 6, isoform 7, isoform 8 and isoform 9 do not bind IL15. Signal transduction involves STAT3, STAT5, STAT6, JAK2 (By similarity) and SYK.,PTM:A soluble form (sIL-15RA) arises from proteolytic shedding of the membrane-anchored receptor. The cleavage involves ADAM17/TACE (By similarity). It also binds IL-15 and thus interferes with IL-15 binding to the membrane receptor.,PTM:N-glycosylated and O-glycosylated.,PTM:Phosphorylated by activated SYK.,similarity:Contains 1 Sushi (CCP/SCR) domain.,subcellular location:Isoform 6, isoform 7, isoform 8 and isoform 9 are associated with endoplasmic reticulum, Golgi and cytoplasmic vesicles, but not with the nuclear membrane.,subcellular location:Mainly found associated with the nuclear membrane.,subunit:The interleukin-15 receptor IL15R is a heterotrimer of IL15RA, IL2RB and IL2RG. IL15RA also self-associates (By similarity). Interacts with SYK.,tissue specificity:Isoform 1, isoform 3, isoform 4, isoform 5, isoform 6, isoform 7, isoform 8 and isoform 9 are widely expressed. Expressed in fetal brain with higher expression in the hippocampus and cerebellum than in cortex and thalamus. Higher levels of soluble sIL-15RA form in comparison with membrane-bound forms is present in all brain structures.,

研究领域 (Research Area)


Cytokine-cytokine receptor interaction;Jak_STAT;Intestinal immune network for IgA production;

图片 (Image Data)




Western blot analysis of lysates from COLO cells, using IL15RA Antibody. The lane on the right is blocked with the synthesized peptide.


产品名称: IL-15R α Rabbit Polyclonal Antibody
产品货号: APRab12502



Western blot analysis of the lysates from HeLa cells using IL15RA antibody.



Western Blot analysis of COLO cells using IL-15R α Polyclonal Antibody



Western blot analysis of various lysis using IL-15R α Polyclonal Antibody. Secondary antibody was diluted at 1:20000

注意事项 (Note)

For research use only .