

产品名称: IL-17RB Rabbit Polyclonal Antibody
产品货号: AP Rab12509



产品概述 (Summary)

产品名称 (Production Name)	IL-17RB Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	IL17RB Interleukin-17 receptor B (IL-17 receptor B) (IL-17RB) (Cytokine receptor-like
别名 (Alternative Names)	4) (IL-17 receptor homolog 1) (IL-17Rh1) (IL17Rh1) (Interleukin-17B receptor) (IL-17B receptor)
基因 ID (Gene ID)	55540.0
蛋白 ID (SwissProt ID)	Q9NRM6.Synthesized peptide derived from IL-17RB at AA range: 251-300

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	55kDa

产品名称: IL-17RB Rabbit Polyclonal Antibody
产品货号: AP Rab12509




研究背景 (Background)

The protein encoded by this gene is a cytokine receptor. This receptor specifically binds to IL17B and IL17E, but does not bind to IL17 and IL17C. This receptor has been shown to mediate the activation of NF-kappaB and the production of IL8 induced by IL17E. The expression of the rat counterpart of this gene was found to be significantly up-regulated during intestinal inflammation, which suggested the immunoregulatory activity of this receptor. [provided by RefSeq, Jul 2008],function:Receptor for the proinflammatory cytokines IL17B and IL17E. May play a role in controlling the growth and/or differentiation of hematopoietic cells.,tissue specificity:Expressed in several endocrine tissues, mostly in fetal and adult liver, kidney, pancreas, testis, colon, brain and small intestine; not detected in peripheral blood leukocytes, lymphoid organs, and most cell lines.,

研究领域 (Research Area)

Cytokine-cytokine receptor interaction;

图片 (Image Data)



Western Blot analysis of HEPG2 3T3 cells using IL-17RB Polyclonal Antibody diluted at 1:1000. Secondary antibody was diluted at 1:20000

注意事项 (Note)

For research use only .