

产品名称: I κ B- β Rabbit Polyclonal Antibody
产品货号: AP Rab12806



产品概述 (Summary)

产品名称 (Production Name)	I κ B- β Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC
种属反应性 (Reactivity)	Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	NFKB1B NFKB1B; IKBB; TRIP9; NF-kappa-B inhibitor beta; NF-kappa-B1B; I-kappa-B-beta; I κ B-B; I κ B-beta; IkappaBbeta; Thyroid receptor-interacting protein 9; TR-interacting protein 9; TRIP-9
别名 (Alternative Names)	
基因 ID (Gene ID)	4793.0
蛋白 ID (SwissProt ID)	Q15653.The antiserum was produced against synthesized peptide derived from human IkappaB-beta. AA range:4-53

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:50-1:300
蛋白分子量 (Molecular Weight)	37kDa

产品名称: I κ B- β Rabbit Polyclonal Antibody
产品货号: AP Rab12806




研究背景 (Background)

The protein encoded by this gene belongs to the NF-kappa-B inhibitor family, which inhibit NF-kappa-B by complexing with, and trapping it in the cytoplasm. Phosphorylation of serine residues on these proteins by kinases marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B, which translocates to the nucleus to function as a transcription factor. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2011],function:Inhibits NF-kappa-B by complexing with and trapping it in the cytoplasm. However, the unphosphorylated form resynthesized after cell stimulation is able to bind NF-kappa-B allowing its transport to the nucleus and protecting it to further IKBA-dependent inactivation. Association with inhibitor kappa B-interacting NKIRAS1 and NKIRAS2 prevent its phosphorylation rendering it more resistant to degradation, explaining its slower degradation.,PTM:Phosphorylated; followed by degradation. Interaction with NKIRAS1 and NKIRAS2 probably prevents phosphorylation.,similarity:Belongs to the NF-kappa-B inhibitor family.,similarity:Contains 6 ANK repeats.,subunit:Interacts with THRB (via ligand-binding domain). Interacts with RELA and REL. Interacts with COMMD1 and inhibitor kappa B-interacting Ras-like NKIRAS1 and NKIRAS2.,tissue specificity:Expressed in all tissues examined.,

研究领域 (Research Area)


Chemokine;NOD-like receptor;RIG-I-like receptor;Cytosolic DNA-sensing pathway;T_Cell_Receptor;B_Cell_Antigen;Neurotrophin;Adipocytokine;

图片 (Image Data)




Western blot analysis of lysates from NIH/3T3 cells, treated with TNF 20ng/ml 30', using IkappaB-beta Antibody. The lane on the right is blocked with the synthesized peptide.

产品名称: IκB-β Rabbit Polyclonal Antibody
产品货号: APRab12806



Western blot analysis of the lysates from HT-29 cells using IκB-β antibody.



Western Blot analysis of NIH-3T3 cells using IκB-β Polyclonal Antibody

注意事项 (Note)

For research use only .