

产品名称: LATS1/2 Rabbit Polyclonal Antibody
产品货号: APRab13234



产品概述 (Summary)

产品名称 (Production Name)	LATS1/2 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	LATS1 WARTS
别名 (Alternative Names)	Serine/threonine-protein kinase LATS1 (EC 2.7.11.1) (Large tumor suppressor homolog 1) (WARTS protein kinase) (h-warts)
基因 ID (Gene ID)	9113.0
蛋白 ID (SwissProt ID)	O95835.Synthesized peptide derived from human LATS1/2 AA range: 1050-1130

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:50-1:300,ELISA 1:2000-1:20000
蛋白分子量 (Molecular Weight)	140kDa

产品名称: LATS1/2 Rabbit Polyclonal Antibody
产品货号: AP Rab13234




研究背景 (Background)

The protein encoded by this gene is a putative serine/threonine kinase that localizes to the mitotic apparatus and complexes with cell cycle controller CDC2 kinase in early mitosis. The protein is phosphorylated in a cell-cycle dependent manner, with late prophase phosphorylation remaining through metaphase. The N-terminal region of the protein binds CDC2 to form a complex showing reduced H1 histone kinase activity, indicating a role as a negative regulator of CDC2/cyclin A. In addition, the C-terminal kinase domain binds to its own N-terminal region, suggesting potential negative regulation through interference with complex formation via intramolecular binding. Biochemical and genetic data suggest a role as a tumor suppressor. This is supported by studies in knockout mice showing development of soft-tissue sarcomas, ovarian stromal cell tumors and a high sensitivity to carcinogenic treatment. catalytic activity: ATP + a protein = ADP + a phosphoprotein., cofactor: Magnesium., function: Tumor suppressor which plays a critical role in maintenance of ploidy through its actions in both mitotic progression and the G1 tetraploidy checkpoint. Negatively regulates G2/M transition by down-regulating CDC2 kinase activity. Involved in the control of p53 expression. Affects cytokinesis by regulating actin polymerization through negative modulation of LIMK1. May also play a role in endocrine function., PTM: Autophosphorylated and phosphorylated during M-phase of the cell cycle. Phosphorylated by STK3 at Ser-909 and Thr-1079, which results in its activation. Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family., similarity: Contains 1 AGC-kinase C-terminal domain., similarity: Contains 1 protein kinase domain., similarity: Contains 1 UBA domain., subcellular location: Localizes to the centrosomes throughout interphase but migrates to the mitotic apparatus, including spindle pole bodies, mitotic spindle, and midbody, during mitosis., subunit: Complexes with CDC2 in early mitosis. LATS1-associated CDC2 has no mitotic cyclin partner and no apparent kinase activity. Binds phosphorylated ZYX, locating this protein to the mitotic spindle and suggesting a role for actin regulatory proteins during mitosis. Binds to and colocalizes with LIMK1 at the actomyosin contractile ring during cytokinesis., tissue specificity: Expressed in all adult tissues examined except for lung and kidney.,

研究领域 (Research Area)

图片 (Image Data)

产品名称: LATS1/2 Rabbit Polyclonal Antibody
产品货号: APRab13234



Western blot analysis of various lysate, antibody was diluted at 1000. Secondary antibody was diluted at 1:20000

注意事项 (Note)

For research use only .