

产品名称: LIR-1 Rabbit Polyclonal Antibody
产品货号: AP Rab13332



产品概述 (Summary)

产品名称 (Production Name)	LIR-1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	LILRB1 LILRB1; ILT2; LIR1; MIR7; Leukocyte immunoglobulin-like receptor subfamily B member 1; LIR-1; Leukocyte immunoglobulin-like receptor 1; CD85 antigen-like family member J; Immunoglobulin-like transcript 2; ILT-2; Monocyte/macrophage immunoglobulin-like receptor 7; MIR-7; CD85j
别名 (Alternative Names)	
基因 ID (Gene ID)	10859.0
蛋白 ID (SwissProt ID)	Q8NHL6.The antiserum was produced against synthesized peptide derived from the N-terminal region of human LILRB1. AA range:21-70

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:50-1:300
----------------------	--------------------------------

产品名称: LIR-1 Rabbit Polyclonal Antibody
产品货号: AP Rab13332



蛋白分子量 (Molecular Weight) 72kDa


研究背景 (Background)

This gene is a member of the leukocyte immunoglobulin-like receptor (LIR) family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008], domain: Contains 4 copies of a cytoplasmic motif that is referred to as the immunoreceptor tyrosine-based inhibitor motif (ITIM). This motif is involved in modulation of cellular responses. The phosphorylated ITIM motif can bind the SH2 domain of several SH2-containing phosphatases., function: Receptor for class I MHC antigens. Recognizes a broad spectrum of HLA-A, HLA-B, HLA-C and HLA-G alleles. Receptor for H301/UL18, a human cytomegalovirus class I MHC homolog. Ligand binding results in inhibitory signals and down-regulation of the immune response. Engagement of LILRB1 present on natural killer cells or T-cells by class I MHC molecules protects the target cells from lysis. Interaction with HLA-B or HLA-E leads to inhibition of the signal triggered by FCER1A and inhibits serotonin release. Inhibits FCGR1A-mediated phosphorylation of cellular proteins and mobilization of intracellular calcium ions., PTM: Phosphorylated on tyrosine residues. Dephosphorylated by PTPN6., similarity: Contains 4 Ig-like C2-type (immunoglobulin-like) domains., subunit: Binds PTPN6 when phosphorylated. Binds FCER1A and FCGR1A., tissue specificity: Expressed predominantly on B-cells and monocytes, and at lower levels on dendritic cells. Detected on a low percentage of T-cells and natural killer (NK) cells.,

研究领域 (Research Area)

Immunology

图片 (Image Data)



Western Blot analysis of SKOV3 cells using LIR-1 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

产品名称: LIR-1 Rabbit Polyclonal Antibody
产品货号: APRab13332



注意事项 (Note)

For research use only .