

产品名称: LMX1B Rabbit Polyclonal Antibody
产品货号: APRab13365



产品概述 (Summary)

产品名称 (Production Name)	LMX1B Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	LMX1B
别名 (Alternative Names)	LMX1B; LIM homeobox transcription factor 1-beta; LIM/homeobox protein 1.2; LMX-1.2; LIM/homeobox protein LMX1B
基因 ID (Gene ID)	4010.0
蛋白 ID (SwissProt ID)	O60663.The antiserum was produced against synthesized peptide derived from human LMX1B. AA range:126-175

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	40kDa

产品名称: LMX1B Rabbit Polyclonal Antibody
产品货号: AP Rab13365




研究背景 (Background)

LIM homeobox transcription factor 1 beta(LMX1B) Homo sapiens This gene encodes a member of LIM-homeodomain family of proteins containing two N-terminal zinc-binding LIM domains, 1 homeodomain, and a C-terminal glutamine-rich domain. It functions as a transcription factor, and is essential for the normal development of dorsal limb structures, the glomerular basement membrane, the anterior segment of the eye, and dopaminergic and serotonergic neurons. Mutations in this gene are associated with nail-patella syndrome. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2010],disease:Defects in LMX1B are the cause of nail-patella syndrome (NPS) [MIM:161200]; also known as Onychoosteodysplasia. NPS is a disease that cause abnormal skeletal patterning and renal dysplasia.,function:Essential for the specification of dorsal limb fate at both the zeugopodal and autopodal levels.,similarity:Contains 1 homeobox DNA-binding domain.,similarity:Contains 1 LIM zinc-binding domain.,similarity:Contains 2 LIM zinc-binding domains.,tissue specificity:Expressed in most tissues. Highest levels in testis, thyroid, duodenum, skeletal muscle, and pancreatic islets.,

研究领域 (Research Area)

图片 (Image Data)



注意事项 (Note)

For research use only .