

产品名称: MAD1 Rabbit Polyclonal Antibody
产品货号: APRab13552



产品概述 (Summary)

产品名称 (Production Name)	MAD1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	MAD1L1 MAD1L1; MAD1; TXBP181; Mitotic spindle assembly checkpoint protein
别名 (Alternative Names)	MAD1; Mitotic arrest deficient 1-like protein 1; MAD1-like protein 1; Mitotic checkpoint MAD1 protein homolog; HsMAD1; hMAD1; Tax-binding protein 181
基因 ID (Gene ID)	8379.0
蛋白 ID (SwissProt ID)	Q9Y6D9.The antiserum was produced against synthesized peptide derived from human MAD1. AA range:394-443

产品应用 (Application)

稀释比 (Dilution Ratio)	ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000
----------------------	--

产品名称: MAD1 Rabbit Polyclonal Antibody
产品货号: APRab13552



蛋白分子量 (Molecular Weight)


研究背景 (Background)

MAD1L1 is a component of the mitotic spindle-assembly checkpoint that prevents the onset of anaphase until all chromosome are properly aligned at the metaphase plate. MAD1L1 functions as a homodimer and interacts with MAD2L1. MAD1L1 may play a role in cell cycle control and tumor suppression. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2015],disease:Defects in MAD1L1 are involved in the development and/or progression of various types of cancer.,function:Component of the spindle-assembly checkpoint that prevents the onset of anaphase until all chromosomes are properly aligned at the metaphase plate. May recruit MAD2L1 to unattached kinetochores. Has a role in the correct positioning of the septum. Required for anchoring MAD2L1 to the nuclear periphery.,induction:Increased by TP53.,PTM:Phosphorylated; by BUB1. Become hyperphosphorylated in late S through M phases or after mitotic spindle damage. Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the MAD1 family.,subcellular location:From the beginning to the end of mitosis, it is seen to move from a diffusely nuclear distribution to the centrosome, to the spindle midzone and finally to the midbody.,subunit:Homodimer. Heterodimerizes with MAD2L1 in order to form a tetrameric MAD1L1-MAD2L1 core complex. Perturbation of the original MAD1L1-MAD2L1 structure by the spindle checkpoint may decrease MAD2L1 affinity for MAD1L1. CDC20 can compete with MAD1L1 for MAD2L1 binding, until the attachment and/or tension dampen the checkpoint signal, preventing further release of MAD2L1 on to CDC20. Also able to interact with the BUB1/BUB3 complex and the viral Tax protein. Interacts with TPR.,tissue specificity:Expressed weakly at G0/G1 and highly at late S and G2/M phase.,

研究领域 (Research Area)

Cell_Cycle_G1S;Cell_Cycle_G2M_DNA;

图片 (Image Data)



Immunofluorescence analysis of NIH/3T3 cells, using MAD1 Antibody. The picture on the right is blocked with the synthesized peptide.

产品名称: MAD1 Rabbit Polyclonal Antibody
产品货号: APRab13552



注意事项 (Note)

For research use only .